
DETERMINANTS OF NET RETURNS TO AGRO FORESTRY IN THE HUMID RAIN FOREST BELT OF NIGERIA.

Obasi, P. C.; Okparadim, G. I. and Henri-Ukoha, A.

Department of Agricultural Economics,
Federal University of Technology, Owerri
P. M. B. 1526, Owerri, Nigeria

ABSTRACT: *The study analyzed the determinants of net returns to agro forestry in the humid rainforest belt of Nigeria. The multi-stage random sampling technique was used in selecting the sample. The sample size comprised 120 agro forestry practitioners who were selected from the list provided by the staff of the Imo state Agricultural Development Programme (ADP). The results of the ordinary least squares multiple regression analysis showed that farm size, years of farming experience, tree crop density, educational attainment, extension contact, type of soil fertility replacement materials used, and farmers age are the major determinants of net returns to agro forestry practice in the state. It was suggested that co-operative farming and communal system of land use could significantly enhance participation in agro forestry in Imo State, Nigeria.*

KEYWORDS: Determinants, Net Returns, Agro Forestry, Humid Rainforest, Nigeria

INTRODUCTION

Agricultural production till date remains the mainstay of the Nigerian economy. Over the years the sector has witnessed tremendous decline in its contribution to national development. This is evident in the high incidence of poverty among the rural populace. The reason for this widespread poverty in Sub Saharan Africa including Nigeria is the destruction of the natural forest leading to environmental degradation and reduced productivity. Besides, small farmers usually farm on degraded lands with complex and diverse farming systems (Owese, 2009). Consequently, population pressures, deforestation and bush burning, farming and other forms of land use (Ekwebalam and Onyenwotu, 1989) are the bane of Nigerian Agriculture. Regrettably, the recent global economic crisis has made it increasingly difficult for families to generate enough resources required for sustainable economic growth without exerting much pressure on natural resources to generate more income and produce more food for household consumption (Akinyemi et al., 2009). The forest area of Nigeria has diminished from above 60 million hectares in the 1990's to the current value of about 9.6 million hectares (NEST, 2004). Loss of forest biodiversity can result from fragmentation, overharvesting of plant and animal species and intensification of arable farmlands. For instance, the land devoted to food crops alone rose from 7.6million hectares in 1970 to 35.5 million hectares in 1995 (Agboola, 1987; FORMECU, 1995). The implication is that more forest lands are subjected to ever increasing pressure by subsistence farmers and herders. This quest for maximum food production for the masses has exerted a negative impact on some of the forest cover through bush clearing and burning. Agro forestry as a concept is the deliberate incorporation of trees and woody species of plants into other types of agricultural activities (Rahman et al, 2011). It is a concept of combining crops, animals and trees

on the same piece of land as maximum land use practice (Beetz, 2002). The technique offers solution to land shortage, poverty, food insecurity and environmental degradation (Lipper, 2002). Again, agro forestry techniques are designed to provide tree and other crop products and at the same time protect, conserve, diversify and sustain vital economic, environmental and natural resources (Kings, 1987; Baumer, 1990; Hawkins et al., 1990). The place of agro forestry in the productivity, sustainability, adaptability, prevention of soil erosion, poverty alleviation and social stability protection has been widely researched (Akpan et al., 2009; Cubbage et al, 2012; Udawatta and Godsey, 2010; Kio, 2000; Henri-Ukoha et al, 2010). However, there is limited information on the determinants of net returns from agro forestry in the study area. In an attempt to close the gap, the study set out to achieve the following objectives:

Objectives of the study

The broad objective of the study was to evaluate the determinants of net returns to agro forestry in the humid rain forest belt of Nigeria with a view to developing strategies for sustainable ecological conservation in Nigeria. The specific objectives are:

- (i) to examine and identify the socio-economic characteristics of agro forestry practitioners in Imo State, Nigeria
- (ii) to ascertain the level of participation of households and communities in agro forestry in the area
- (iii) to identify the types of agro forestry practiced, and the species and varieties of crops planted in the area
- (iv) to determine and isolate the main determinants of net returns to agro forestry in the area.

MATERIALS AND METHODS

Imo state is located in the humid south-eastern region of Nigeria with a total land mass of 7,689 square kilometers (UNTC, 1998). The state has a population of 3,934,899 people (NPC, 2006). The occupation of the people is mainly agriculture. The system of land use in the state can be grouped into forest land, wood land and agricultural land. The state is made up of twenty seven (27) local government areas divided into three (3) major agricultural zones for administrative convenience. The three agricultural zones are Owerri, Orlu and Okigwe respectively. The multi-stage stratified random sampling technique was used in selecting the respondents. Firstly, two local government areas were randomly selected from each of the three agricultural zones of the state. This gave a total of six local government areas. In the second stage, two autonomous communities were randomly selected from each of the six selected local government areas, giving a total of twelve communities. Thirdly, ten agro forestry practitioners were selected randomly from each of the twelve communities thus giving a sample size of one hundred and twenty (120) farmers. The list of agro forestry practitioners in the area was supplied by staff of the Imo state agricultural development programme (ADP). Data for the study were collected mainly from a primary source. The primary data were obtained using structured questionnaire which was administered to the selected farmers through oral interview. Data were analyzed using descriptive statistical tools such as means, frequencies and percentages. The ordinary least squares multiple linear regression was used to analyze the factors affecting the returns to agro forestry in the state. The estimated model is expressed as:

$$Y_t = b_0 + b_1X_{1t} + b_2X_{2t} + \dots + b_8X_{8t} + e_t \quad \dots \dots \dots \text{eqn.(1)}$$

Where,

Y_t = Returns to agro forestry production (₦)

X_{1t} = Age of farmers (years)

X_{2t} = Farm size (Ha)

X_{3t} = Years of farming experience (years)

X_{4t} = Tree crop density (number of stands)

X_{5t} = Educational attainment (years)

X_{6t} = Household size (persons)

X_{7t} = Extension contact (dummy 1= yes, 0 = otherwise)

X_{8t} = Type of soil fertility enhancing material used (dummy, 1= organic manure; 0= otherwise)

e = Error term

Different functional forms of the model were estimated and the one that gave the best fit was selected as the lead equation and used for further analysis of the data. The viability of agro forestry practice in the state was measured using benefit cost ratio (BCR) and net present value (NPV). These are specified below:

$$\text{Benefit Cost Ratio (BCR)} = \frac{\text{total discounted benefits}}{\text{total discounted costs}} \dots\dots\dots \text{eqn.(2)}$$

$$\text{Benefit Cost Ratio} = \frac{\sum_{i=1}^n B_n(1+i)^n}{\sum_{i=1}^n C_n(1+i)^n} \dots\dots\dots \text{Eqn.(3)}$$

$$\text{NPV} = \text{cash flow}_0 + \text{cashflow}_1[1/(1+i)^1] + \text{cashflow}_2[1/(1+i)^2] + \dots + \text{cash flow}_n[1/(1+i)^n] \dots\dots\dots \text{eqn.(4)}$$

Since cashflow_0 is not affected by the variability of the discount factor, it is moved to the other side of the equation.

$$\text{NPV} = \text{cash flow} \left[\sum_{t=1}^n \frac{1}{(1+i)^t} \right] \dots\dots\dots \text{eqn.(5)}$$

The cash flow is the annual equivalent value (AEV) that is being calculated.

$$\text{Cash flow} = \frac{\text{NPV}}{\sum_{t=1}^n \frac{1}{(1+i)^t}} \dots\dots\dots \text{eqn.(6)}$$

Simplifying eqn. (5), we obtain the series of annual equivalent values of the investment as

$$\text{AEV} = \sum_{t=1}^n \frac{1}{(1+i)^t} = \frac{1}{i} - \frac{1}{i(1+i)^n} \dots\dots\dots \text{eqn.(7)}$$

Where,

BC R = Benefit Cost Ratio

NPV = Net Present Value

Cash flow_n = Net income or net loss for the year “n”

Cash flow₁ = Net income from the first full year of production.

i = Discount rate or the opportunity cost of investing.

n = Number of years included in the budget.

Similarly, the net returns to agro forestry in the state was estimated using the eqns. (8 to 10)

$$TR_i = \sum_{i=1}^n P_q Q \dots\dots\dots \text{eqn.}(8)$$

$$TVC_i = \sum_{i=1}^n P_i X_i \dots\dots\dots \text{eqn.}(9)$$

Where,

TR = Total Revenue generated

TVC = Total Variable costs

P_q = Unit price of output

Q = Quantity of output produced

X = Quantity of the variable input

$$\text{Net Returns} = TR_{ij} - (TVC + TFC) \dots\dots\dots \text{eqn.}(10)$$

TFC = Total fixed costs

RESULTS AND DISCUSSION

Socio-economic characteristics of respondents.

Table 1. Distribution of agro forestry practitioners according to socio-economic characteristics

Socio-economic characteristics	Frequency	Relative Frequency
Gender		
Male	95	79.16
Female	25	20.83
Age (Years)		
≤ 30	2	1.67
31 – 40	6	5.00
41 – 50	62	51.67
51 – 60	28	23.33
61 – 70	12	10
71 – 80	7	5.83
81 and above	3	2.50

Mean age = 51years		
Marital status		
Single	4	3.33
Married	110	91.67
Widowed	6	5.00
Household size (persons)		
≤ 5	27	22.50
6 – 10	68	56.67
11 – 15	20	16.67
16 and above	5	4.17
Mean household size = 8persons		
Educational attainment (years)		
No formal education	8	6.67
6 – 10	28	33.33
11 -15	72	60
16 and above	12	10
Mean years of formal education = 7		
Farm size (Ha)		
≤ 1.0	34	28.33
1.1 – 2.0	62	51.67
2.1 – 3.0	16	13.33
3.1 – 4.0	4	3.33
4.1 – 5.0	2	1.67
5.1 and above	2	1.67
Mean farm size = 1.4 hectares		
Years of farming experience		
≤ 10	40	33.33
11 – 20	64	53.33
21 – 30	8	6.67
31 – 40	6	5.00
41 and above	2	1.67
Mean years of farming experience = 14		
Total	120	

Source: Survey data, 2010

Table 1 shows the distribution of respondents according to socio-economic characteristics. The Table shows that 79.16% of the sampled agro forestry practitioners are males while 20.83% are females.

This suggests the involvement of more males in agro forestry practice in the area. With respect to age, the Table shows that the majority (51.7%) are within the age bracket of 41 – 50 years. The mean age was 51 years. This may indicate that agro forestry in the state is practiced more by active middle aged men and women who could adopt new techniques in agro forestry farming. Apart from increase in labour supply, respondents within the productive age bracket are likely to adopt innovation more than the aged farmers (Obasi et al, 2012). The Table also shows that 91.67% of the respondents are married, 3.33% are single, while 5.00% are widows. It could be deduced from the analysis that married people do readily engage in agro forestry practice than unmarried people. Furthermore, the Table showed that about 67.6% of the respondents have household sizes that range between 6 – 10 persons. The mean household size was 8 persons. With regard to educational attainment, the study showed that the farmers had a minimum of seven years of formal education, while the highest percentage had between 7 – 12 years of education. The level of education of a person not only increases his farm productivity but also enhances his ability to understand and evaluate new production technologies (Henri-Ukoha et al, 2011). In addition to the above, the study further show that 53.3% of the respondents had spent between 11 – 12 years in agro forestry farming. The mean years of agro forestry farming was 13.8 years. Similarly, 51.67% of the respondents have farm sizes that range between 1.1 – 2.0 hectares. The mean farm size was 1.67 hectares. This implies that agro forestry in Imo State is practiced on a small scale.

Participation in agro forestry

Table 2. Distribution of respondents by level of participation in agro forestry

Forestry activities	High	Moderate	Low/mild	No participation
Nursery management	11 (09)	31 (26)	28 (23)	52 (43)
Home gardens	36 (30)	40 (33)	19 (16)	25 (21)
Alley cropping	01 (01)	07 (06)	19 (16)	91 (76)
Multipurpose trees	13 (11)	25 (21)	23 (19)	59 (49)
Crops/trees in rotation	16 (13)	31 (26)	24 (20)	49 (41)
Improved fallow	13 (11)	29 (24)	25 (21)	53 (44)
Shelterbelts/wind brakes	10 (08)	24 (20)	31 (26)	55 (46)
Trees and hedgerows	13 (11)	13 (11)	13 (11)	79 (66)
Plantation crops	24 (20)	41 (34)	17 (14)	40 (33)
Trees as live fences	18 (15)	29 (24)	24 (20)	49 (41)
Farm woodlots	18 (15)	40 (33)	16 (13)	48 (40)
n = 120				

Figures in parenthesis are percentages

Source: Field survey data, 2010.

Table 2 shows the distribution of respondents by level of participation in agro forestry. The Table shows that (9%) of the respondents actively participated in the management of nurseries, 26% moderately participated in nursery management, 23% did not actively participate, while 43% do not manage nurseries. Similarly, 30% highly participated in home gardening, 33% moderately grew home gardens, and 16% did not actively grow home gardens, while 21% do not grow home gardens at

all. The Table also shows that 1% highly participated in alley cropping, 6% moderately participated in alley cropping, and 16% mildly participated in alley cropping, while 76% did not participate in alley cropping. Furthermore, 11% of the respondents actively grew multipurpose trees, 21% moderately grew multipurpose trees, and 19% did not actively grow multipurpose trees, while 49% did not grow multipurpose trees at all. Also, 13% actively practiced crops and trees in rotation, 26% moderately practiced crops and trees in rotation; 20% did not practice crops and trees in rotation, while 41% did not practice crops and trees in rotation at all. The Table also shows that 8% actively grew shelterbelts and windbreaks, 20% moderately grew shelterbelts and windbreaks, 11% did not actively grow shelterbelts and windbreaks, while 66% did not grow shelterbelts and windbreaks at all. Similarly, 20% actively grew plantation crops, 34% moderately grew plantation crops, 14% did not actively grow plantation crops, while 33% did not grow plantation crops at all. Furthermore, 15% actively grew trees as live fences, 24% moderately grew trees as live fences, 20% did not actively grow trees as live fences, while 41% did not grow trees as live fences at all. Finally, 15% were highly involved in growing farm woodlots, 33% are moderately involved in growing farm woodlots, 13% did not actively grow farm woodlots, while 40% were not involved in growing farm woodlots at all.

Types of agro forestry systems practiced

Table 3. Distribution of respondents according to types of agro forestry practiced

Type of agro forestry	Frequency	Relative percentage
Plantation crops	35	29.17
Farm wood lots	12	10.00
Fruit trees and crops in combination		
Trees as live fences	77	64.17
Trees and hedgerows	11	9.17
Multistory home gardening	05	4.17
Dispersed multipurpose trees	13	10.83
Improved fallow	02	1.66
Wildlife agro forestry	10	8.33
Amenity planting	02	1.66
Alley farming	04	3.33
	01	0.83

Source: Field survey data, 2010

Table 3 shows the distribution of respondents according to types of agro forestry practiced. The Table shows that fruit trees and crops in combination (64.17%), and plantation trees (29.17%) were cultivated due to their roles in food security, the variety of crops obtained, logs and timber derived, and income generation. Other agro forestry practices in the area included trees and hedgerows(4.12%), live fences (9.17%), farm woodlot (10.00%), and multistory home gardening (10.83%) which were used because of their roles in enhancement of aesthetic beauty, control of wind and water erosion, supply of fodder, fuel wood and soil fertility maintenance.

Types of soil fertility replacement materials used by Farmers**Distribution of respondents according to type of soil fertility replacement materials used in Agro forestry practice**

Soil fertility replacement material	Frequency	Relative frequency
Animal waste	20	16.67
Green manures (mulching)	34	28.33
Household and kitchen waste	14	11.67
Inorganic fertilizers	74	61.67

Source: Field survey data, 2010

Table 4 shows the distribution of respondents according to type of soil fertility replacement materials used in Agro forestry practice. The Table shows that 16.67% of the respondents used animal waste as soil fertility enhancement material, 28.33% used green manures, 11.67% used household and kitchen waste, while 61.67% used inorganic fertilizers. The result suggests that majority of the respondents (61.67%) use inorganic fertilizer as soil fertility replacement material.

Determinants of net returns to agro forestry production in Imo State**Results of the multiple regression analysis on the determinants of net returns to agro forestry in Imo State.**

Variable	Coefficients	T-ratio
Age (X ₁)	-0.0743 (0.0614)	1.2101
	0.0937	
Farm size (X ₂)	(0.0213)	4.3991*
	0.0742	
Years of experience in agro forestry (X ₃)	(0.0169)	4.3905*
	0.0318	
Tree crop density practiced (X ₄)	(0.0107)	2.9719*
	0.0664	
Years of formal education (X ₅)	(0.0185)	3.5892*
	0.0903	
Household size (X ₆)	(0.0817)	1.1053
	0.0885	
Extension contact (X ₇)	(0.0307)	2.8827*
	0.0685	
Fertility replacement material (X ₈)	(0.0329)	2.2313**
	73.4447	
Constant	(0.0329)	
R ² = 0.7142		
F-Value = 34.3365, n = 120		

Figures in parentheses are standard errors

* significant at 1% ** significant at 5%

Source: Field survey data, 2010

Following Olayide and Heady (1982), the double log function was chosen as the lead equation and used for further analysis of the data. The results of the analysis show that Farm size (X_2), years of farming experience (X_3), tree crop density (X_4), years of formal education (X_5), extension contact (X_6) and soil fertility replacement material used (X_8) are statistically significant within 1% to 5% levels and directly related to net returns to agro forestry. This suggests that an increase in the levels used of these inputs will significantly enhance net returns to agro forestry in the area.

Table 6: Discounted Benefits and Costs from agro forestry practice in Imo State

Year	Gross revenue	Discount factor (18%)	Present worth	Gross costs	Discount factor (18%)	Present worth
1	116895	0.848	99126.960	54197	0.848	45959.056
2	142529	0.718	102335.822	68583	0.718	49242.594
3	172729	0.609	105191.961	66013	0.609	40201.917
4	148300	0.516	76522.800	59648	0.516	30778.368
5	138033	0.437	60320.421	54750	0.437	23925.750
6	146935	0.370	54365.950	57329	0.370	21211.730
7	151841	0.314	47678.074	60789	0.314	19087.746
8	153518	0.266	40835.788	83963	0.266	22334.158
9	142770	0.226	32266.020	53887	0.266	12178.462
10	114553	0.191	21879	41646	0.191	7954.386
TOTAL			640523.419			272874.167

$$\text{BC Ratio} = \frac{\text{present worth of benefits}}{\text{present worth of costs}} = \frac{640523.419}{272874.167} \quad \text{BC Ratio} = 2.347$$

Since BC Ratio is greater than 1, the agro forestry enterprise is viable.

Table 7: Discounted cash flow from agro forestry practice in Imo State

Year	Gross revenue (1)	Gross costs (2)	Cash flows (1-2)	Discount factor (18%)	Present worth
1	116895	54197	62698	0.848	53167.904
2	142529	68583	73946	0.609	64990.044
3	172729	66013	106716	0.609	64990.044
4	148300	59648	88652	0.516	45744.432
5	138033	54750	83283	0.437	36394.671
6	146935	57329	89606	0.370	33154.220
7	151841	60789	91052	0.314	28590.328
8	153518	83963	69555	0.266	18501.630
9	142770	53887	88883	0.226	20087.558
10	114553	41646	72907	0.191	13932.528
TOTAL					₦ 367,656.543

NPV = ~~₦~~367,656.543

Since NPV is positive, the agro-forestry enterprise is economically viable.

Applying equations (6 and 7) to Table 7, we obtain

Where NPV = 367656.543, i = 18% (market interest rate), t = 10 years (planning horizon)

$$\text{Cash flow} = \sum_{t=1}^n \frac{1}{(1+i)^t} = \frac{1}{0.18} - \frac{1}{0.18(1+0.18)^{10}} = 4.49$$

$$\text{Therefore, AEV} = \frac{NPV}{4.49} = \frac{367656.543}{4.49} \quad \text{AEV} = \del{₦}81,883.417$$

The result indicates that the series of cash flows expected from the agro forestry practice have the same net present value as an annuity that pays ~~₦~~81,883.417 per year. This does not reflect the time it started generating positive cash flows. Therefore the practice of agro forestry in the state is economically viable.

Table 8: Net returns to agro forestry practice in Imo State.

Item	Value (₦ 1years)
Total revenue (TR)	142810.34
Variable cost (VC)	
Seed planting materials	17573.83
Chemical fertilizer	14505.25
Organic fertilizer	1042.00
Labour	22618.30
Agro-chemical	2745.50
Total variable cost (TVC)	58484.90
Fixed cost (FC)	
Depreciation on capital items	4574.69
Land rent	6425.71
Total fixed costs (TFC)	11000.40
Total costs (TC)	69485.30
Net revenue (TR-TC)	73325.04

BCR = 2.055

USA1 dollar = NGN145

Source: Field survey data, 2008

Table 8 shows a positive net returns of ₦73,325.04. The benefit-cost ratio is 2.10 implying that for every one naira (₦1.00) spent in the agro forestry business, one naira ten kobo is returned (₦1.10k). This result indicates the viability of agro forestry practices in Imo state.

CONCLUSION

Based on the estimated BCR and NPV, we conclude that agro forestry practice in the humid rainforest belt of Nigeria is economically viable and should be invested upon.

References

- Agboola, S.A. (1987). *Agricultural Atlas of Nigeria*. Oxford University Press, Ibadan, P.148
- Akinyemi, G.O., Adebago, C.A., Tayo, O.S., Akinyemi, I.G and Ige, P.O (2009). *Effective Environmental Management: A recipe for Sustainable Food Production in Nigeria*, Pp. 775-778
- Akpan, U.F., Nsien, I.B and Udoh, G.E (2009). *Tackling Economic Financial Meltdown through a Traditional Agricultural Practise: Agroforestry*. Proceedings of the 43rd Annual Conference of the Agricultural Society of Nigeria, Abuja, 2006. Pp. 779-780
- Baumer, M. (1990). *The Potential role of Agro-forestry in combating desertification and Environmental degradation with specific reference to Africa*. Technical Centre for Agricultural and Rural Corporation (ACP-EEC Lome Convention. The Netherlands.
- Beetz, A. (2002). *Agroforestry Overview Horticultural System Guide*. Attra Publication.
- Cabbage, F.W,V. Glenn, J. P. Mueller, D. Robison, R. Myers, J. M. Luginbuhl and R. Myers (2012). *Early tree growth, crop yields and estimated returns for an agroforestry trial in Goldsboro, North Carolina*. . *Agroforestry Systems: An International Journal in corporation with ICRAF*. Vol. 71, No. 3.
- Ekwebalam, S.A and Onyewotu, L.O (1989). *Management for environmental forestry*. In: F.A. Akinsamni (ed). *Proceedings of National Workshop on “Strategies for Environmental Forestry Management in Nigeria”*.
- FORMECU (1995). *An Asseement of Land Use and Vegetaation Changes in Nigeria, 1978-1995*, Geomatis, Ontario, Canada.
- Hawkins, R., Sembiring, H., Lubis, D and Suwardjo (1990). *The Potential of Alley Cropping in the Uplands of East and Central Java*. Upland Agriculture Conservation Project (UACP), Agency for Agricultural Research and Development, Department of Agriculture, Salatiga, Indonesia.
- Henri-Ukoha, A., Orebiyi, J.S., Ohajianya, D.O., Nwosu, F.O., Nwaiwu, I.U., Ben-Chendo, G.N (2010). *Economics of dry season vegetable production by women farmers in Owerri West Local Government Area of Imo State, South East Nigeria: A case study of telfaria occidentalis*. *International Journal of Agriculture and Rural Development*; Vol. 12, Pp.32-36
- Henri-Ukoha A., J. S. Orebiyi, P. C. Obasi, N. N. Oguoma, D. O. Ohajianya, U. C. Ibekwe and I. I. Ukoha (2011). *Determinants of Loan acquisition from the Financial Institutions by Small-scale Farmers in Ohafia Agricultural zone of Abia State, South-east Nigeria*. *Journal of Development and Agricultural Economics* 3(2), 69-74.
- Kings, K.F.S (1987). *The History of Agroforestry* In: Stepphre, H.A and Nair, P.K.R (eds). *Agroforestry a decade of development*, Nairobi, Kenya. (ICRAF, Pp1-11.
- Kio, P.R.O (2000). *Forestry and Sustainable Agricultural Development*. National Workshop on Agriculture and Rural Development in Nigeria. In: Owonubi, J.J; Oguntala, A.B and Soladoye, M.O. (eds).
- Lipper, L (2002). *Forest degradation and Food Security Unasylva* 202, vol. 53, 1997.

- NEST (2004), Nigeria Climate Change: Executive Summary of five multi-sector-sector surveys on Nigeria's vulnerability and adaptation to climate. Canada-Nigeria Climate Change Project.
- NPC (2006). National Population Commission. Nigerian Agriculture Magazine Vol. 4 No.3 Pp. 22 August/September 2000
- Obasi, P. C.;Okparadim,G.I.;and Henri-Ukoha, A. (2012): Economics of Agroforestry in Imo State, Nigeria. International Journal of Agricultural and Food Science. Vol. 2, No. 1. Pp. 7 – 13.
- Owese, T.O., Odebode, A.V., and Amadi, J.O (2009). Prospects of Agroforestry as a veritable tool in mitigating food insecurity and poverty in Nigeria. Proceedings of the Agricultural Society of Nigeria, Abuja, 2009. Pp. 781-783
- Rahman, S. A., F. D. Paras, S. R. Khan, A. Imtiaj, K. M. Farhana, M. M. Toy, M. B. Akhand and T. Sunderland (2011). Initiatives of tropical Agro-forestry to Sustainable Agriculture: A case study of Capasia Village, Northern Bangladesh. Journal of Horticulture and Forestry Vol. 3(4), Pp. 115-121
- Udawatta R. P, Godsey L. D (2010). Agroforestry comes of age: putting science into practice. Agroforest System 79(1): Pp.1–4
- United Nation program of Technical Cooperation U.N.P.TC (1998). Demographic and health data collection and analysis in Nigeria. Project No. NIR98P40. New York, Ny UN.
- Yudelman, M. (1987). Prospects for Agricultural Development in Sub Sahara Africa. Occasional Paper (Winrock International Institute for Agricultural Development) April, Little Rock, Arkansas.