

EFFECT OF CO-ADMINISTRATION OF CHLOROQUINE PHOSPHATE AND CEFUROXIME AXETIL ON SERUM LIPIDS AND LIPID PEROXIDATION IN ALBINO WISTAR RATS

Etim O. E*, Akpan E. J. and Akpanabiatu M. I.

Department of Biochemistry, Faculty of Basic Medical Sciences, University of Uyo, Nigeria

ABSTRACT: *There is scanty information on the response of biochemical system to the co-administration of anti-malarial drugs and antibiotics. This was what informed the need to undertake the present study. The effects of co-administration of chloroquine (CQ) and cefuroxime (CE) on serum lipids and lipid peroxidation in albino Wistar rats was investigated. Standard reagent kits was used for the assay. The study included assay on the effect of the co-administration of CQ and CE on high density lipoprotein cholesterol (HDL-C), total cholesterol (CHOL), triglycerides (TG), low density lipoprotein cholesterol (LDL-C) and lipid peroxidation. The result revealed that the level of HDL-C was not affected by the co-administration. Those of TC, TG and LDL-C increased significantly ($p < 0.05$) above the value obtained for the control group. Malondialdehyde (MDA) level increased significantly ($p < 0.05$) in group treated with CQ in combination with CE. The present study has revealed that co-administration of chloroquine and cefuroxime tends to raise TC, TG, LDL-C and MDA levels. The result reveals that co-administration of chloroquine and cefuroxime may adversely affect the metabolism of lipids.*

KEYWORDS: Chloroquine, Cefuroxime, Serum Lipids, Lipid Peroxidation.

INTRODUCTION

It is well established in the literature that bacterial infections can coexist with severe malaria, especially in the tropics. Rosenthal *et al.*, (2010) advocated that despite appropriate anti-malarial therapy, these patients should receive broad spectrum antibiotic. Moreover in malaria endemic areas, children often carry falciparum malaria asymptotically, so malaria is over-diagnosed at the expense of other infectious conditions (Maitland, 2010). A study carried out to access the coexistence of bacterial infection with *Plasmodium falciparum* revealed that blood slide positive for *Plasmodium falciparum* had also invasive bacterial disease, and improved diagnosis and treatment of invasive bacterial disease along with malarial disease are needed to reduce childhood mortality.

It is also commonplace in recent times that at the end of malarial therapy, symptoms which may or may not be of malarial origin seem to persist. Reasons for these persistent symptoms may not be known until proper medical laboratory tests are conducted. However in an attempt to solve the problem, certain antibiotics are administered alongside anti-malarials. This co-administration ultimately eradicates the infections (bacterial or parasitic). It is not well established or known whether the drug combination is safe to the biochemical system or not.

There is scanty information on the response of the biochemical system to the co-administration of anti-malarial and antibiotics. Therefore this study was imperative as the effects of the co-administration of an anti-malarial and antibiotic on the albino Wistar rats will be investigated.

MATERIALS AND METHODS

Materials

Antimalarial: Chloroquine phosphate (250 mg)

Chloroquine phosphate (250 mg) is manufactured by Evans Medical PLC (RC 1161), Km 32, Lagos – Badagry Expressway, Agbara Industrial Estate, Agbara, Ogun State, Nigeria.

Antibiotic: Cefuroxime axetil tablets (500 mg)

Cefuroxime axetil tablets (500 mg) is manufactured by Okasa Pharma PVT Ltd, L-2 Additional MIDC Area, Satara 415004- India. It is manufactured for CIPLA Ltd India under loan licence.

It marketed by Evans Medical Plc, Agbara Industrial Estate, Agbara, Ogun State, Nigeria

Methods

Experimental design and treatment of animals

A total of twenty four (24) albino Wistar rats (male and female) were weighed with average weight of 151g and divided into four (4) different groups of six (6) per group and put into rat cages. Group 1 rats were given normal rat pellets and water. Group 2 rats were treated with 4.17 mg/kg body weight (bw) of chloroquine (CQ). Group 3 rats were administered concomitantly with 4.17 mg/kg body weight (bw) of chloroquine and 8.33 mg/kg bw of cefuroxime (CE). Group 4 rats were treated with 8.33 mg/kg bw of cefuroxime (CE) only.

Assay of biochemical parameters:

Estimation of high density lipoprotein-Cholesterol (HDL-C) Level was carried out according to the method of Lopes Virella *et al*, 1997. Low density lipoproteins (LDL and VLDL) and chylomicrons fractions are precipitated quantitatively by the addition of phosphotungstic acid in the presence of magnesium ions. After centrifugation, the cholesterol concentration in the HDL (high density lipoprotein) fraction, which remains in the supernatant, is determined. Estimation of Serum Cholesterol (CHOL) level was carried out according to the method of Allain *et al*, 1974; Roeschlaw *et al*, 1974. The concentration of cholesterol is determined after enzymatic hydrolysis and oxidation. The indicator, quinoneimine is formed from hydrogen peroxide and 4-aminoantipyrine in the presence of phenol and peroxidase (Trinder, 1969). Estimation of Serum Triglycerides (TRIGS) level was carried out according to the method of Tietz, 1990. The triglycerides are determined after enzymatic hydrolysis with lipases. The indicator is a quinoneimine formed from hydrogen peroxide, 4 – aminophenazone and 4 – chlorophenol under the catalytic influence of peroxidase. Low density lipoprotein-Cholesterol (LDL-C) was carried out according to the method of Friedewald *et al.*, 1972.

LDL Cholesterol = Total cholesterol – Triglycerides – HDL Cholesterol

Assessment of Lipid Peroxidation (LPO) was carried out according to the method of Esterbauer *et al*, 1991.

Assessment of lipid peroxidation is based on the reaction of a chromogenic reagent, N-methyl-2-phenylindole (RI) with malondialdehyde (MDA) and 4-hydroxyalkenals at 45 °C. One molecule of either MDA or 4-hydroxyalkenals reacts with 2 molecules of R1 to yield a

stable chromophore with maximal absorbance at 586 nm. For simultaneous estimation of MDA concentration and 4-hydroxyalkenals, one must use the procedure utilising methanesulfonic acid (MSA) as the acid solvent. The procedure in which HCl is used will only detect MDA, since the 4-hydroxyalkenals does not form a chromophore with reagent R1 under those conditions.

Statistical Analyses:

Data obtained were analysed using appropriate statistical methods such SPSS statistics, analysis of variances (ANOVA). Results were considered significant at ($p < 0.05$).

RESULTS AND DISCUSSION

Results

Table 1: Serum Lipid Profile of albino Wistar rats Co-Administered WITH 4.17 mg/kg bw of CQ and 8.33 mg/kg bw of CE

Treatment	TC (mmol/L)	TG(mmol/L)	HDL-C (mmol/L)	LDL-C (mmol/L)
CTR	1.84±0.05	5.90±0.16	0.58±0.01	0.30±0.09
CQ	2.28±0.16*	6.91±0.24*	0.54±0.01	0.39±0.12*
CQ+CE	1.79±0.08**	6.45±0.09*	0.57±0.04	0.57±0.09*
CE	1.66±0.072**	6.49±0.06*	0.54±0.01	0.75±0.09*

CQ = chloroquine, CE = cefuroxime

* significantly different from control (CTR) ($P < 0.05$), ** Significantly different within group, number of rats (n) = 6.

Table 2: Effects of the Co-Administration of 4.17 mg/kg bw of CQ and 8.33 mg/kg bw of CE on Lipid Peroxidation in albino Wistar Rats (Assessed in terms of MDA Level)

Treatment	MDA (µmol)
CTR	4.76±0.36
CQ	3.93±0.41**
CQ+CE	5.97±0.30*
CE	4.28±0.22**

CQ = chloroquine, CE = cefuroxime

* significantly different from control (CTR) ($P < 0.05$), ** Significantly different within group, number of rats (n) = 6.

DISCUSSION

High prevalence of malaria and typhoid fever in the tropics has made co-infections common. However, the actual and precise underlying mechanism to explain the association between malaria and salmonellosis is still uncertain (Uneke, 2008).

The treatment of malaria and typhoid co-infection is a common phenomenon in many parts of Africa (Nsutebu and Ndumbe, 2001). This is because malaria and typhoid fever have been associated with poverty and underdevelopment with significant morbidity and mortality. An association between malaria and typhoid fever was first described in the medical literature in the middle of the 19th century, and was named typhomalarial fever by the United States Army (Smith, 1982).

Rosenthal *et al*, (2010) advocated that despite appropriate anti-malarial therapy, these patients should receive broad spectrum antibiotics. Moreover, in malaria endemic areas, children often carry falciparum malaria asymptotically, so malaria is over diagnosed at the expense of other infectious conditions.

Serum lipid profile in the treated Wistar rats is presented on Table 1. Cholesterol is a fatty substance found in blood, bile and brain tissue. It serves as a precursor for the synthesis of bile acids, steroids, and vitamin D. The determination of serum cholesterol is a major aid in the diagnosis and classification of lipemia (Tietz, 1990). Other conditions such as hepatic thyroid diseases influence cholesterol level (Jacobs and Van Demark, 1960). They act as functional and structural components of biomembranes and form insulation to allow nerve conduction and prevent heat loss.

High density lipoproteins (HDL) are one of the major classes of plasma lipoproteins. They are composed of a number of heterogenous particles, including cholesterol and vary with respect to size and content of lipid and apolipoprotein. HDL serves to remove cholesterol from the peripheral cells to the liver, where the cholesterol is converted to bile acids and excreted into the intestine. The apparent fixed level of HDL in the present study literally implies that HDL level was not affected by the drugs treatment. Rahilly (2011) advocated that for a fixed level of HDL, the risk of heart disease increases three-folds as LDL varies from low to high.

Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction. In the human body, high levels of triglycerides in the blood stream have been linked to atherosclerosis (hardening of the arteries), and by extension the risk of the heart disease and stroke (AHA, 2009).

Total cholesterol (TC), triglycerides (TG) and low density lipoprotein cholesterol (LDL-C) increase significantly ($P < 0.05$) as shown in Table 1 when the group treated with CQ was compared to control. Triglycerides (TG) and LDL-C also increased significantly ($p < 0.05$) when the group treated with CQ+CE and the group treated with CE alone were compared to the control, an indication of increased risk of heart disease.

An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. The importance of HDL-C as a risk factor for CHD is now recognized (NIHCDCS, 1992).

Data from the Landmark Framingham Heart Study revealed that, for a given level of LDL, the risk of heart disease increases 10-folds as the HDL varies from the high to low. On the converse, however, for a fixed level of HDL, the risk increases 3-folds as LDL varies from low to high (Rahilly, 2011).

Malondialdehyde reacting substances and serum in treated Wistar rats were shown on Table 2. Lipid peroxidation, a known mechanism of cellular injury is a chain reaction initiated by oxidant radicals. It is an index of oxidative stress in cells and tissues. The free radical oxidation of lipid molecules proceeds from the fatty acylmethylene group adjacent to a double bond (Frankel, 1980) to form lipid hydroperoxides which are capable of further reactions to yield various classes of peroxides. Lipid peroxides are unstable and decompose to form series of complex products including reactive carbonyl compounds. Polyunsaturated fatty acids on oxidation decompose to malondialdehyde (MDA) and 4 – hydroxyalkenals (HAE). The levels of these products in serum or tissues have been widely accepted as a measure of lipid peroxidation (Esterbauer *et al*, 1991).

Biological membranes which contain phospholipids rich in polyunsaturated fatty acids are continuously being oxidised by superoxides, hydroperoxyl and peroxy radicals emanating from exposures to radiations and oxidative reactions involving endogenous and exogenous compounds in the body (Droge, 2002).

The toxicity of lipid hydroperoxides to animals is best illustrated by the lethal phenotype glutathione peroxidase 4 (GPx4) knock out mice. These animals do not survive, past embryonic day 8, indicating that removal of lipid hydroperoxides is essential for mammalian life (Muller *et al.*, 2007).

Data from the present study, revealed a significant increase ($p < 0.05$) in MDA concentration in the group treated with CQ+CE compared to the control. The result is an indication of potential toxicity to membrane by the drug combination though the level of toxicity noticed does not seem to trigger the antioxidant defense system observed in this study. These systems include antioxidant enzymes, transport and storage proteins, etc. Oxidative stress which is associated with cellular injury in many pathological conditions occur when there is an imbalance between free radical generation and radical scavenging systems (Hensley *et al*, 2000).

REFERENCES

- Allain, C.C., Poon, L.S., Chan, C.S., Richmond, W.S. and Fu, P.C (1974). Enzymatic Determination of Total Serum Cholesterol. *Clinical Chemistry*. 20: 470 – 475.
- American Heart Association (AHA). (2009). Your Triglyceride Levels. What your cholesterol levels mean. <http://www.americanheart.org>.
- Droge, W. (2002). Free radicals in the Physiological Control of Cell Function. *Physiological Review.*, 82:47 – 52.
- Esterbauer, H., Schaur, R.J. and Zollner, H. (1991). Chemistry and Biochemistry of 4-Hydroxyalkenal, Malondialdehyde and Related Aldehydes. *Free Radical Biology and Medicine.*, 11: 81 – 128
- Frankel, E.N. (1980). Lipid Peroxidation. *Progress of Lipid Research.*, 19:1 – 22.
- Friedwald, W.T., Levy, R. I. and Fredrickson, D. S. (1972). Estimation of the Concentration of Low Density Lipoprotein Cholesterol in Plasma, without Use of the Preparative Ultracentrifuge. *Clinical Chemistry.*, 18: 499.
- Henley, K., Robinson, K.A., Gabbita, S.P., Salsman, S. and Floyd, R.A. (2000). Reactive Oxygen Species, Cell Signaling, and Cell Injury. *Free Radical Biology and Medicine.*, 28: 1456 – 1461.

- Jacobs, N. J. and VanDemark, P. J. (1960). The Purification and Properties of the Alpha Glycerophosphate Oxidizing Enzyme of *Streptococcus faecalis* loci. *Archives of Biochemistry and Biophysics.*, 88: 250 – 255.
- Lopes-Virella, M.F., Stone, P., Ellis, S. and Colwell, J. A. (1977). Cholesterol Determination in High Density Lipoproteins Separated by Three Different Methods. *Clinical Chemistry.*, 23(5): 882 – 884.
- Maitland, K. (2010). Antimicrobials in Children Admitted to Hospital in Malaria Endemic Areas. *Biomedical Journal.*, 340: 1818.
- Muller, F. L., Lustgarten, M. S., Jang, Y., Richardson, A. and Van Remmen, H. (2007). Trends in Oxidative Aging Theories. *Free Radical Biology and Medicine.*, 43: 477 – 503.
- National Institutes of Health Consensus Development (NIHCDCS). (1992). Conference Statement: Triglyceride, High Density Lipoprotein and Coronary Heart Diseases. Washington D.C.
- Nsutebu, E. and Ndumbe, P. (2001). The Widal Test for Typhoid Fever: Is it Useful? Oguce, S., Omokhodion, S.I., Adeyemo., A.A. and Olumeses, P.E. (2002). Low Plasma Bicarbonate Predicts Poor Outcome of Cerebral Malaria in Nigeria Children. *West African Journal Medicine.*, 21(4): 45 – 46.
- Rahilly, C. (2011). Relation between High-density Lipoprotein Cholesterol and Survival to Age 85 Years in Men (from the VA normative aging study). *American Journal of Cardiology.*, 21: 296 – 300.
- Roeschlaw, P., Bernt, E. and Guber, J. W. (1974). *Clinical Chemistry and Clinical Biochemistry.*, 12: 403.
- Rosenthal, N., Vinciguerra, M., Santini, M.P., Claycomb, W.C. and Ladurner, A.G. (2010). Local IGF-1 Isoform Protects Cardiomyocytes from Hypertrophic and Oxidative Stresses via SirT1 Activity. *AGING-US.* 2: 43 – 62.
- Smith, D. C. (1982). The Rise and Fall of Typhomalarial Fever. In: Origins. *Journal of Historical Medical and Allied Science.*, 37: 182 – 220
- Tietz, N. W. (1990). *Clinical Guide to Laboratory Tests.* 2nd ed, Philadelphia, U.S.A: W.B. Saunders Co. pp. 554 – 556.
- Uneke, C. J. (2008). Concurrent Malaria and Typhoid Fever in the Tropics: The Diagnostic Challenges and Public Health Implications. *Journal of Vector Borne Diseases.*, 45: 133 – 142.