

## HISTO-MORPHOMETRIC EFFECTS OF SILDENAFIL CITRATE ON THE TESTIS OF NORMOGLYCAEMIC AND HYPERGLYCAEMIC ADULT WISTAR RATS

J. E. Ataman<sup>1</sup> and C. L. Sakpa<sup>2</sup>

<sup>1</sup>Department of Anatomy, School of Basic Medical Sciences, College of Medical Sciences, University of Benin, Benin-City, Nigeria.

<sup>2</sup>Department of Anatomy, College of Medicine of the University of Lagos, Nigeria.

**ABSTRACT:** *Diabetes mellitus is a known cause of infertility and erectile dysfunction in men. Sildenafil citrate, used in the management of the latter, has been implicated of toxic role on the testis. This paradoxical impression is significant in diabetic men with erectile dysfunction that are of reproductive age. Histo-morphometric effects of sildenafil citrate was thus, investigated for eight weeks on the testes of twenty-five adult Wistar rats, weighed before and after the experiment and categorized into four treatment groups and one control group of five rats per group (n=5). The control group (A) received feed mash and water ad libitum. Treatment groups B and C (normoglycaemic) received low dose (1mg/kg body weight) and high dose (2mg/kg body weight) of sildenafil citrate, respectively. Groups D and E (hyperglycaemic) were treated with low dose (1 mg/kg body weight) and high dose (2 mg/kg body weight) of sildenafil citrate, respectively. Blood samples were collected for hormonal assay and the testes were excised and processed for morphological changes. The results showed significant ( $P<0.05$ ) loss in body and testicular weight in the hyperglycaemic groups, but insignificant ( $P>0.05$ ) body weight gain in the normoglycaemic groups and insignificant ( $P>0.05$ ) epididymal weight difference in all the treatment groups, compared to control. The hormone assay showed significant ( $P<0.05$ ) difference in the levels of FSH, LH, testosterone and oestrogen in the hyperglycaemic groups compared to control. Testicular and epididymal tissues revealed mild distortions in hyperglycaemic treatments but with dose dependent improvement, while the changes in the normoglycaemic treatments were essentially non-remarkable. The results suggest relative safety of sildenafil citrate in normoglycaemic and hyperglycaemic states, with dose-dependent beneficial testicular effects in the latter condition.*

**KEYWORDS:** Sildenafil Citrate, Normoglycaemia, Hyperglycaemia, Testis, Wistar rats

## INTRODUCTION

A very well known endocrine and metabolic disorder that affects males and females across diverse age range in whom sexual dysfunction can occur is diabetes mellitus (Ballester *et al.*, 2004; Akef *et al.*, 2012). In the male, infertility and impotence are two threatening complications associated with diabetes mellitus (Zhao *et al.*, 2011; Ataman and Osinubi, 2014; Oyelade *et al.*, 2016). Thus, diabetes might be co-morbidity in erectile dysfunction (Brown *et al.*, 2005; El-Sakka *et al.*, 2008). There is increasing evidence that diabetes is closely associated with male reproductive dysfunctions with observed pathological changes in Leydig cells, interstitial connective tissues and seminiferous tubules of the testis (Nascimento Silva *et al.*, 2014; Ataman and Osinubi, 2017). Compared with non-diabetic people, male diabetic patients show an increasing incidence of erectile dysfunction and infertility with erectile

dysfunction occurring about ten years earlier in the diabetics than in the non-diabetic counterparts (Kiskac *et al.*, 2015). Experimental diabetic animals tend to suffer from testicular dysfunction such as reduced sperm count, low serum testosterone levels and decreased fertility (Mallick *et al.*, 2010; Ataman and Osinubi, 2017).

Erectile dysfunction has been identified as a possible complication in about 50 % of diabetic men (Spollett, 1999). Poor glycaemic control has been related to the pathogenesis of diabetes mellitus which results in complications as vasculopathy, neuropathy and myopathy (Kiskac *et al.*, 2015; Anwar *et al.*, 2017), which interferes with the normal stimulation of noradrenergic and noncholinergic nerves in the pelvic parasympathetic plexuses that cause release of nitric oxide across the neuromuscular junction of the penile arteries and cavernosa smooth muscles (Spolletti, 1999; McCullough, 2000). With diabetes mellitus as a known risk factor in erectile dysfunction, decreased endothelial levels of nitric oxide synthase has been implicated in its pathogenesis and mediated phosphorylation of this enzyme reportedly, promotes penile erection (Hurt *et al.*, 2002). Current treatment modality for it with tremendous acceptance is seen in the role of drug formulations as sildenafil citrate, a phosphodiesterase- 5 (PDE-5) inhibitor which helps in potentiating the levels of nitric oxide by binding to the PDE-5 enzyme, preventing PDE-5 breakdown of cyclic guanosine monophosphate (cGMP) through competitive inhibition. Nitric oxide causes increase in cGMP which results in the relaxation of penile smooth muscles and increased cavernosa blood flow to promote penile erection (McCullough, 2002; Montague *et al.*, 2005).

The role of sildenafil on testicular morphology and functions has been associated with mixed submissions, either emphasizing on its efficacy or its possibly deleterious effects on the testis. While it has been reported (Yildiz *et al.*, 2011) that sildenafil citrate served a cytoprotective role via reduction in oxidative stress in a study with compromised vascular supply to the testis, it was also reported as not without some morphological distortions to some of the testicular tissues following graded dosage. Its low dose in another comparative study (Suriyakumari *et al.*, 2015) elicited adverse effects with accumulated/depleted metals and cytoarchitectural distortions on interstitial space and spermatogenic cells, capable of causing impotency which led to their querying the ability of sildenafil to bring about a reversal effect. On the other hand, sildenafil citrate has also been reported to have positive effects on spermatogenesis, sperm production, and semen quality (Beheshtian *et al.*, 2008). These are equivocal submissions on the role of sildenafil on the testis.

It is known that poor glycaemic control in men of reproductive age leads to attendant infertility challenge and vulnerability to erectile dysfunction (Brown *et al.*, 2005; Agbaje *et al.*, 2007), that might require long term treatment with sildenafil citrate (Vardi and Nini, 2007; Schouten *et al.*, 2010). An important issue for consideration is does sildenafil ameliorate infertility challenge posed by diabetes in its effects on testicular functions or does it worsen it? In this study, the morphometric and morphological effects of graded dose application of sildenafil citrate on normal and induced hyperglycaemia in Wistar rats is evaluated to provide a clue.

## **MATERIALS AND METHOD**

### **Drug Preparation and Administration**

The drug, sildenafil citrate is manufactured by HAB Pharmaceuticals Limited was obtained from Thelver Pharmacy, Benin City. Drug preparation was done by diluting a tablet of 100mg in 50mls of distilled water to obtain 2mg/ml, and administration was 1mg/kg body weight and 2mg/kg body weight.

### **Experimental animals**

Twenty-five Wistar rats bred at the Animal house of the Department of Anatomy, University of Benin, Benin City were used for this study under ethical approval. The animals were kept in standard environmental conditions of humidity of  $65 \pm 5\%$ , room temperature of between 25 to 26°C and 12:12 hours of day and night photoperiodicity. Experimental procedures involving the animals and their care were conducted in conformity with International and Institutional guidelines for the care of laboratory animals in Biomedical Research, as promulgated by Canadian Council of Animal care (Canadian Council of Animal Care, 1985). Further, the animal experimental models used were in conformity to the guiding principles for research involving animals as recommended by the Declaration of Helsinki and the Guiding Principles in the care and use of Animals (American Physiological Society, 2002).

They were categorized into four treatment groups and one control group of five rats per group. The duration of the experiment was two months. The control group (A) was fed on feed mash only and water throughout the period. The four treatment groups (B, C, D and E) received feed mash, water and sildenafil citrate but groups D and E were induced with diabetes using 50mg/kg body weight of streptozotocin manufactured by Sigma Aldrich Company, USA. Confirmation of diabetes at 250mg% was done with accu check glucometer and strips before the administration of sildenafil Citrate. The treatment regimen is shown in Table 1. The body weights of the animals were measured at the beginning and end of the experiment and recorded.

After eight weeks of administration, the animals were sacrificed by cervical dislocation. Blood samples were collected for hormone assay using ELISA kits in line with the guiding principles of the tests in DRG Diagnostics User's Manual. The abdomen was excised to retrieve the testes and the epididymis which were processed through prior 24 hours fixation in bouin's fluid. The fixed tissues were passed through ascending grades of alcohol and the dehydrated tissues were cleared in xylene before infiltration with molten paraffin wax and thereafter allowed to cool and solidify. The embedded tissues were trimmed and mounted on a wooden chauck and sectioned at 5 microns using a rotary microtome. Staining with hematoxylin and eosin was done (Drury and Wallington, 1980) and the stained slides were mounted with Canada balsam and coverslipped for microscopy which was carried out at x400 magnification.

### **Data Analysis**

Data were presented as Mean  $\pm$  SEM. Means separation and significant differences between the means (Duncan, 1957) were determined at ( $P < 0.05$ ) using ANOVA.

**Table 1: Showing Experimental Groups and the Treatment Regimen**

GROUPS	TREATMENT REGIMEN
<b>A</b>	Received Feed mash and water ad libitum
<b>B</b>	Feed mash and low dose of sildenafil citrate (1mg/kg body weight) in normoglycaemic rats.
<b>C</b>	Feed mash and high dose of sildenafil citrate (2mg/kg body weight) in normoglycaemic rats.
<b>D</b>	Hyperglycaemic rats: received water ad-libitum, feed mash and low dose of sildenafil citrate (1mg/kg body weight)
<b>E</b>	Hyperglycaemic rats: received water ad-libitum, feed mash and high dose of sildenafil citrate (2mg/kg body weight)

## RESULTS

From the results of the effects of treatments on the mean weight values of experimental rats as shown in Table 1, the initial mean body weight was  $246.00 \pm 9.14$  g, while the final mean body weight value was  $257.00 \pm 7.68$  g for group A (Control) animals. In group B, the initial mean body weight value changed from  $189.00 \pm 6.40$  g to the final mean body weight value of  $209 \pm 6.78$  g. In Group C, the initial mean body weight value was  $208.00 \pm 9.82$  g while the final mean weight value was  $219.00 \pm 7.31$  g. In Group D, the initial mean body weight value of  $229.00 \pm 5.34$  g dropped to final mean weight value of  $214.00 \pm 5.34$  g. In Group E, the initial mean body weight value of  $227.00 \pm 6.44$  g became  $223.00 \pm 0.02$  g at the end of the experiment. The findings revealed weight gain in groups B and C, but not significantly ( $P > 0.05$ ) different from control. Conversely, there was significant ( $P < 0.05$ ) weight loss in the hyperglycaemic groups D and E compared to control.

The results on mean testicular weight of the rats post-sacrifice was significant ( $P < 0.05$ ) in groups D ( $1.22 \pm 0.02$  g) and E ( $0.88 \pm 0.32$  g) compared to control ( $1.41 \pm 0.18$  g) but there was however no significant difference ( $P > 0.05$ ) between the mean testicular weight of the other treatments and the control. Also the results of the mean epididymal weight revealed no significant difference ( $P > 0.05$ ) between that of the treatment groups and the control (Table 2).

Table 3 showed the hormonal profile of the experimental groups. From the results, the values of FSH ( $0.08 \pm 0.32$  IU/ml), LH ( $0.91 \pm 0.01$  IU/ml) and testosterone ( $1.02 \pm 0.12$  ng/ml) in the hyperglycaemic group were significantly lesser ( $P < 0.05$ ) in group D compared to control where the values were  $0.13 \pm 0.44$  IU/ml,  $1.23 \pm 0.1$  IU/ml and  $2.72 \pm 1.31$  ng/ml, respectively. The oestrogen value was unaffected by sildenafil treatment in groups B ( $13.13 \pm 1.24$  ng/ml) and C ( $14.25 \pm 1.32$  ng/ml), but insignificantly ( $P > 0.05$ ) higher in the hyperglycaemic groups D ( $17.36 \pm 2.04$  ng/ml) and E ( $15.34 \pm 1.50$  ng/ml) compared to control ( $14.22 \pm 0.03$  ng/ml). The prolactin values for groups B ( $0.82 \pm 0.42$  ng/ml), C ( $1.01 \pm 0.02$  ng/ml), D ( $0.83 \pm 0.11$  ng/ml) and E ( $0.77 \pm 0.23$  ng/ml) were not significantly different ( $P > 0.05$ ) from control ( $0.94 \pm 0.55$  ng/ml). Also, there was no significant difference ( $P > 0.05$ ) in the values of progesterone between the

treatment groups B ( $2.35 \pm 0.12$  ng/ml), C ( $2.02 \pm 0.02$  ng/ml), D ( $1.89 \pm 0.45$  ng/ml) and E ( $1.99 \pm 0.26$  ng/ml) and the control ( $2.84 \pm 0.33$  ng/ml).

**Table 2: Mean values of the body weight, weight of the testes and epididymis following the administration of sildenafil citrate**

	<b>Group A</b> (mean± S.E.M)	<b>Group B</b> (mean ± S.E.M)	<b>Group C</b> (mean ± S.E.M)	<b>Group D</b> (mean± S.E.M)	<b>Group E</b> (mean± S.E.M)
<b>Initial body Weight</b>	246.00±9.14	189.00±6.40	208.00±9.82	229.00±5.34*	227.00±6.44*
<b>Final body Weight</b>	257.00±7.68	209.00±6.78	219.00±7.13	214.00±5.34*	223.00±9.02*
<b>Weight of Testes</b>	1.41±0.18	1.37±0.09	1.41±0.19	1.22±0.02*	0.88±0.32*
<b>Weight of epididymi s</b>	0.14±0.02	0.12±0.00	0.13±0.01	0.12±0.00	0.11±0.01

Values are Mean ± SEM. Means with asterick remark are significantly different from control: (P<0.05). Horizontal comparisons only

**Table 3: Mean hormonal profile and standard error of mean of control and treatment animals**

Groups	Oestrogen (ng/ml)	FSH (IU/ml)	LH (IU/ml)	Prolactin (ng/ml)	Progesterone (ng/ml)	Testosterone (ng/ml)
A:	14.22±0.03	0.13±0.44	1.23±0.10	0.94±0.55	2.84±0.33	2.72±1.31
B:	13.13±1.24	0.11±0.02	1.07±0.47	0.82±0.42	2.35±0.12	2.14±0.32
C:	14.25±1.32	0.13±0.22	1.22±0.22	1.01±0.02	2.02±0.02	2.36±0.02
D:	17.36±2.04	0.08±0.32*	.91±0.01*	0.83±0.11	1.89±0.45	1.02±0.12*
E:	15.34±1.50	0.10±0.11	1.09±0.13	0.77±0.23	1.99±0.26	2.82±1.20

\* Values are Mean ± SEM. Means with asterick remark are significantly different from control: (p<0.05). Vertical comparisons only

Group A (Control): Received feed mash and water only

Group C: Normoglycaemic with high dose sildenafil

Group D: Hyperglycaemic with low dose sildenafil

Group B: Normoglycaemic with low dose sildenafil

Group E: Hyperglycaemic with high dose sildenafil

FSH: Follicle stimulating hormone

LH: Luteinizing Hormone, Ng/ml: nanogram/milliliters

### Histological findings


The histological outlook of the processed testicular and epididymal specimens from the various experimental groups are shown in Figs. 1-10. The control slides of group A (Figs. 1 & 2) revealed testis with seminiferous tubules in normal sequential maturation, separated by interstitial space and enclosed tunica albuginea while the epididymis showed ducts packed with mature spermatozoa. The normoglycaemic rat testis of group B treated with low dose sildenafil citrate showed apparently normal seminiferous tubules, interstitial vascular hypertrophy, mild congestion and moderate tissue separation while its epididymal tissue had mildly reduced luminal spermatozoa population (Figs. 3 & 4). Group C normoglycaemic rat testis treated with high dose sildenafil citrate similarly had apparently normal seminiferous tubules and moderate tissue separation while its epididymal lumen is enriched with spermatozoa storage but with mild infiltrates of chronic inflammatory cells in the interstitial space (Figs. 5 & 6). Group D hyperglycaemic rat testis treated with low dose sildenafil citrate showed sequential maturation




of sperm cells and its epididymis had mildly reduced luminal spermatozoa population (Figs. 7 & 8). Group E hyperglycaemic rat testis treated with high dose of sildenafil citrate revealed moderate improvement in sequential maturation of spermatozoa and the epididymis showed fairly improved luminal storage of spermatozoa (Figs. 9 & 10).




**Fig 1:** Section of control testis: Seminiferous tubules with normal sequential maturation A, separated by interstitial space B and enclosed tunica albuginea C (H&E x 400)



**Fig 2:** Control rat epididymis showing epithelial lining with columnar cells A, lumen packed with mature spermatozoa B (H&E x 400)




**Fig 3:** Normoglycaemic rat testis treated with low dose sildenafil citrate showing apparently normal seminiferous tubule A, interstitial vascular hypertrophy and mild congestion B and moderate tissue separation C (H&E x 400).




**Fig 4:** Normoglycaemic rat epididymis treated with low dose sildenafil citrate showing mildly reduced luminal spermatozoa population A (H&E x 400).







**Fig 5:** Normoglycaemic rat testis treated with high dose sildenafil citrate showing apparently normal seminiferous tubules A and moderate tissue separation B (H&E x 400).




**Fig 6:** Normoglycaemic rat epididymis treated with high dose sildenafil citrate showing lumen with enriched spermatozoa storage A and mild infiltrates of chronic inflammatory cells in the interstitial space B (H&E x 400).




**Fig 7:** Hyperglycaemic rat testis treated with low dose sildenafil citrate showing normal sequential maturation of sperm cells A (H&E x 400)



**Fig 8:** Hyperglycaemic rat epididymis treated with low dose sildenafil citrate showing mildly reduced luminal spermatozoa population A (H&E x 400)



**Fig 9:** Hyperglycaemic rat testis treated with high dose sildenafil citrate showing moderate improvement in sequential maturation A and medullary interstitium with Leydig cell B (H&E x 400).



**Fig 10:** Hyperglycaemic rat epididymis treated with high dose sildenafil citrate showing fairly increased luminal spermatozoa population A (H&E x 400).

## DISCUSSION

Sildenafil citrate is a known drug of choice in the treatment of erectile dysfunction which is the inability to sustain a satisfactory erection (Boolell *et al.*, 1996; Vardi and Nini, 2007). One of the major causes of erectile dysfunction is diabetes mellitus (Montague *et al.*, 2005) which has also been known to be deleterious to the testis, affecting male fertility (Zhao *et al.*, 2011).



Experimental diabetic animals tend to suffer from testicular dysfunction such as reduced sperm count, low serum testosterone levels and decreased fertility (Mallick *et al.*, 2010).

The result of this study revealed that sildenafil citrate does not significantly affect body, testicular and epididymal weight of treatments compared to control. The observed body, testicular and epididymal weight loss in the hyperglycaemic group is consistent with previous reports (Roy *et al.*, 2013). The normoglycaemic groups treated with low and high doses of sildenafil citrate showed relatively normal testicular and epididymal cytoarchitecture with progressive spermatogenesis in the seminiferous tubules and good spermatogenic storage in the epididymis.

The treatment group with induced hyperglycaemia and low dose sildenafil citrate showed evidence of distorted seminiferous epithelium due to tubular oedema and luminal depletion of spermatozoa which is indicative of impaired spermatogenesis supporting previous report (Suriyakumari *et al.*, 2015). Similar picture was depicted in the lumen of the epididymis of this treatment group with mildly reduced luminal store of spermatozoa. These findings are consistent with earlier report (Khalid, 2009) on significantly reduced sperm count, abnormal morphology and distorted testicular architecture with sildenafil. However, in contrast, the hyperglycaemic group with higher dose of sildenafil citrate in this study showed seminiferous tubules in relatively normal spermatogenesis evidenced with germ cell series in sequential maturation from the basal surface towards the adluminal compartment and matured spermatozoa abutting the lumen. The interstitial compartment is composed of Leydig cells with no cytoarchitectural abnormality. The lumen of the epididymis in contrast to that of the low dose sildenafil citrate hyperglycaemic group is richly enclosed with spermatozoa with no cytoarchitectural distortion.

These results gave the picture of improved spermatogenesis with increasing dose of sildenafil citrate from 1 to 2mg/kg body weight to 2mg/kg. This is supportive of previous reports on the role of sildenafil in apparently compromised testis (Beheshtian *et al.*, 2008; Sivasankaran *et al.*, 2008; Yildiz *et al.*, 2011).

The morphological observations in the testis and epididymis are reflective of the hormonal interplay of the pituitary-hypo-gonadal axis. Impaired hypothalamo-pituitary function with low basal levels of FSH and LH with associated normal or low response to stimulation has been observed as characteristic of diabetic state in animals (Kirchick *et al.*, 1982). The functions of the testes are influenced by gonadotropic hormones produced by the anterior pituitary. Follicle stimulating hormone initiates spermatogenesis and the Luteinizing hormone (LH) results in testosterone release. The presence of both testosterone and follicle-stimulating hormone (FSH) is needed to support spermatogenesis. It has also been shown in animal studies that if testes are exposed to either too high or too low levels of estrogens (such as estradiol; E2), spermatogenesis can be disrupted to such an extent that the animals become infertile (Sierens *et al.*, 2005). Sub-chronic treatment of sildenafil citrate (Viagra) on some enzymatic and non-enzymatic antioxidants in testes and brain of male rats was said to have caused significantly increased MDA levels in testes and significantly reduced brain levels but with observed significant increase in GSH content of testes and brain that suggested therapeutic dose of sildenafil citrate to have elicited modulatory roles by stabilizing/boosting antioxidant defense systems in male rat (Akintunde *et al.*, 2012).

In this study, it was remarkable that impaired testicular spermatogenesis was noted in the hyperglycaemic treatments especially on low dose sildenafil citrate with altered oestrogen and testosterone levels compared to control, while undistorted or insignificantly different levels of FSH, LH and testosterone levels compared to control as noted in the high dose sildenafil citrate-treated hyperglycaemic rats, facilitated spermatogenesis. This finding may be adducible to the possible role of oxidative stress and endothelial dysfunction in induced diabetes (Molnar *et al.*, 2005; Ataman and Osinubi, 2017), which this study considers remediable with the use of higher dose of sildenafil citrate. This position is supported by previous findings (Ayala *et al.*, 2007; Oudot *et al.*, 2009 Mammi *et al.*, 2011; Akef *et al.*, 2012) .

## CONCLUSION

The objective of this study to ascertain the safety of the use of sildenafil citrate, moreso on long term basis in the management of erectile dysfunction in males of reproductive age without compromising fertility is justifiable from these reports as no significantly deleterious effect was noticed in the testicular and epididymal functions following sildenafil administration at the higher dose. However, low dose application did not elicit such as favorable result, consistent with the earlier report (Suriyakumari *et al.*, 2015). Also, the findings from this study did not particularly elicit testicular functional impairment with the use of sildenafil in both normoglycaemic and hyperglycaemic states, but a comparative advantage of possible synergistic effect with its use that is dose dependent. This calls to mind the need to emphasize discretionary use of sildenafil (Smith and Romanelli, 2005) preferably with physician's prescription and monitoring so as to avert its abuse and deleterious effects.

## REFERENCES

- Agbaje, I.M., Rogers, D.A., McVicar, C.M., McClure, N., Atkinson, A.B, Mallidis, C and Lewis, S.E (2007). Insulin-dependent diabetes mellitus: Implications for male reproductive function. *Human Reproduction*, 22(7): 1871-1877.
- Akef, K., Osama, M., Sammah, E. and Safy, G. (2012). Effect of sildenafil on gonadotrophin and sex steroids in fructose induced diabetes in female rats. *Med J Cairo Univ.*, 80(2): 243-252.
- American Physiological Society. (2002). Guiding Principles for Research Involving Animals and Human Beings. *Am J Physiol Regul Integr Comp Physiol* 283: R281 – R283.
- Anwar. Z., Sinha, V., Mitra, S., Mishra, A.K., Ansari, M.H., Bharti, A., Kumar, V. and Nigan, A.K. (2017). Erectile dysfunction: An underestimated presentation in patients with diabetes mellitus. *India Journal of Psychological Medicine*, 39(5): 600-604.
- Ataman, J. E. and Osinubi, A. A. A. (2014). Effects of Streptozotocin-induced Diabetes Mellitus on the Testis of Wistar Rats. *NISEB Journal*, 14(2): 67 –75.
- Ataman, J. E. and Osinubi, A. A. A. (2017). Morphological Evaluation of the effects of Ethanolic leaf- extract of *Newbouldia laevis* (P. Beauv.) on Streptozotocin-induced Gonadotoxicity in Adult Male Wistar Rats. *Zimbabwe Journal of Science & Technology*, 12: 8-18.
- Ayala, J.E., Bracy, D.P., Julien, B.M., Rottman, J.N., Fueger, P.T and Wasserman, D.H (2007). Chronic treatment with sildenafil improves energy balance and insulin action in high fat fed conscious mice. *Diabetes* 56(4): 1025-1033.



- Ballester, J., Munoz, M.C., Dominguez, J., Rigau, T., Guinovart, J.J, Rodriguez-GI, J.E. (2004). Insulin dependent diabetes affects testicular function by FSH- and LH-linked mechanisms. *J Androl.*, 25(5): 706-719.
- Beheshtian, A., Salmasi, A.H, Payabvash, S., Kiumehr, S., Ghazinezami, B., Rahimpour, S., Tavangar, S.M., Dehpour, A.R. (2008). Protective effects of sildenafil administration on testicular torsion/detorsion damage in rats". *World J Urol.*, 26(2):197-202.
- Boolell, M., Allen, M.J., Ballard, S.A., Gepi-Attee, S., Muirhead, G.J., Naylor, A.M., Osterloh, I.H., Gingell, C. (1996). Sildenafil: an orally active type 5 cyclic GMP-specific phosphodiesterase inhibitor for the treatment of penile erectile dysfunction. *Int. J. Impot. Res.* 8 (2): 47–52.
- Brown, J.S., Wessells, H., Chancellor, M.B *et al.*, (2005). Urologic complications of diabetes. *Diabetes care*, 28: 177.
- Canadian Council of Animal Care. (1985). Guide to the Handling and Use of Experimental Animals. NH Publications, Ottawa, USA, vol. 23, Pp: 45 – 47.
- Duncan, D.B. (1957). Multiple Range Test for Correlated and Heteroscedastic Means. *Biometrics* 13: 164 – 176.
- Drury, R.A.B., Wallington, E.A. (1980). Light Microscope and Slide Preparation. Carleton's Histological Technique, 5<sup>th</sup> ed., Oxford University Press, London. Pp.1 – 4.
- do Nascimento Silva, A.A, Oliveira, R.R., de Oliveira, J.S., Nerves, E. (2004). Evaluation of quantitative parameters of Leydig cell in diabetic adult rats. *Acta Scientiarum Biological Sciences*, 36(4): 483-489.
- El-Sakka, A.I., Sayed, H.M., Tayeb, K.A. (2008). Type 2 diabetes-associated androgen alteration in patients with erectile dysfunction. *Int J Androl.*, 31: 602.
- Hurt, K.J., Musicki, B., Palese, M.A., Crone, J.K., Becker, R.E. (2002). Akt- Dependent Phosphorylation of Endothelial Nitric- Oxide Synthase mediates penile erection. 2002. Proceeding of the National Academy of Science, USA. pp. 4061-4066.
- Khalid, G.A. (2009). Effect of long term administration of sildenafil citrate (Viagra) on some sperm characteristics and testes architecture of male rats. *Bas. J. Vet. Res.*, 8(2): 91-103.
- Kirchick, H.J., Keyes, P.L. and Frye, B.E (1982). Restoration of the LH surge and ovulation by insulin in alloxan diabetic immature rats treated with pregnant mare's serum gonadotrophin. *Acta Endocrinologica*, 100: 266-273.
- Kiskac, M., Zorlu, M., Cakirca, M., Buyukaydin, B., Karatoprak, C., Yarovuz, E. (2015). Frequency and determinants of erectile dysfunction in Turkish diabetic men. *Nigerian Journal of Clinical Practice*, 8(2): 209-212.
- Mallick, C., Bera, T.K, Ali, K.M., Chartterjee, K. and Ghosh, D. (2010). Diabetes-induced testicular disorders vis-a vis germ cell apoptosis in abino rat: Remedial effect of hexane fraction of root of *Musa paradisiaca* and leaf of *Coccinia indica*. *Journal of Health Science*, 56 (6): 641-654.
- Mammi, C., Pastore, D., Lombardo, M.F, Ferrelli, F., Caprio, M., Consoli, C., *et al.*, (2011). Sildenafil reduces insulin–resistance in human endothelial cells. *PLoS ONE*, 6(1): e 14542
- McCullough, A.R. (2002). Four year review of sildenafil citrate. *Rev Urol.* 4(3): S24-S38.
- Molnar, J., Yu, S., Mzhavia, N., Pau, C., Chereshevnev, I. and Dansky, H.M. (2005). Diabetes induces endothelial dysfunction but does not increase neointimal formation in high fat diet fed C57BL/6J mice. *Circ. Res.* 96: 1178-1184.

- Montague, D.K., Jarow, J.P., Broderick, G.A., Dmochowski, R.R., Heaton, J.P., Lue, T.F., Milbank, A.J., Nehra, A., Sharlip, I.D. (2005). "Chapter 1: The management of erectile dysfunction: an AUA update". *J. Urol.* 174 (1): 230–239.
- Oudot, A., Behr-Roussel, D., Compagnie, S., Caisey, S., LeCoz, O., Giuliano, F. et al., (2009). Endothelial dysfunction in insulin-resistant rats is associated with oxidative stress and COX pathway dysregulation. *Physiol. Res.* 58: 499-509.
- Oyelade, B.O., Jemilohun, A.C., Aderibigbe, S.A. (2016). Prevalence of erectile dysfunction and possible risk factors among men of South-Western Nigeria: a population based study. *Pan Afr Med J*, 24: 124.
- Roy, S., Rahaman, N., Ahmed, F., Metya, S., Sannigrahi, S. (2013). Naringenin Attenuates Testicular damage, Germ Cell Death and Oxidative Stress in Streptozotocin-induced Diabetic Rats: Naringenin Prevents Diabetic Rat Testicular damage. *J Appl Biomed.*, 11: 195 – 208.
- Schouten, B.W., Bohnen, A.M., Groeneveld, F.P., Dohle, G.R., Thomas, S., Bosch, J.L. (2010). Erectile dysfunction in the community: trends over time in incidence, prevalence, GP consultation and medication use- the Krimpen study: trends in ED. *J Sex Med.*, 7(7): 2547-2553.
- Sierens, J. E., Sneddon, S. F., Collins, F., Millar, M. R., Saunders, P. T. (2005). "Estrogens in Testis Biology". *Annals of the New York Academy of Sciences*, 1061: 65–76.
- Sivasankaran, T.G., Udayakumar, R., Elanchezhiyan., Sabhanayakan, S. (2008). Effect of sildenafil citrate (Viagra) and ethanol on the albino rat testes: A scanning electron microscopic approach. *Cell Biology International*, 32(2): 293-297.
- Smith, K.M., Romanelli, F. (2005). Recreational use and misuse of phosphodiesterase inhibitors. *J Am Pharm Assoc*, (2003) 45(1):63-72.
- Spollett, G.R. (1999). Assessment and Management of erectile dysfunction in men with diabetes. *Diabetes Educ.*, 25: 65-73.
- Suriyakumari, K.V.P., Kumar, U.R., Savitha, D. (2015). Trace element and light microscopic study on testis of Albino rats treated with sildenafil citrate (Caverta). *Int J Anat Res* 3(2): 1168-1172.
- Vardi, M. and Nini, A. (2007). Phosphodiesterase inhibitors for erectile dysfunction in patients with diabetes mellitus" *Cochrane Database Syst Rev* 24(1): CD002187.
- Yıldız, H., Durmus, A.S., Şimşek, H., and Yaman, M. (2011). Protective effect of sildenafil citrate on contralateral testis injury after unilateral testicular torsion/detorsion. *Clinics (Sao Paulo)* 66(1): 137-142.
- Zhao, Y., Tan, Y., Dai, J., Li, B., Guo, L., Cui, J., Wang, G., Shi, X., Zhang, X., Mellen, N., Li, W., Cai, L (2011). Exacerbation of Diabetes- induced Testicular Apoptosis by Zinc Deficiency is most likely Associated with Oxidative Stress, p38 MAPK activation, and p53 Activation in Mice. *Toxicol let.*, 200 (1-2): 100 – 106.