

HISTORICAL RECORDS OF THE LESSEPSIAN MIGRANTS, THE DUSKY SPINEFOOT FISH *SIGANUS LURIDUS* (RUPPELL, 1829) AND THE MARBLED SPINEFOOT *S. RIVULATUS* (FORSSKAL, 1775), IN THE EASTERN COAST OF LIBYA MEDITERRANEAN SEA

Ezalnaser A. Farag Abziew^{*1} and Sayed Mohamed Ali²

¹ Department of Zoology, Faculty of Art and Science, Omar Al-Mukhtar University, Darna, Libya.

² Department of Zoology, Faculty of Science, Omar Al- Mukhtar University, Albaida, Libya.

ABSTRACT: A brief comprehensive account on Lessepsian migration is given in the present study. Earlier publications reported that the dusky spinefoot fish *Siganus luridus* (Ruppell, 1829) and the marbled spinefoot *S. rivulatus* (Forsskal, 1775) were recorded in Libya Mediterranean coastal waters for the first time in 1968 and 1970 in order. The present study however, shows that they were in Libya since the 1950ties: in Tobrok in 1950, in Darna in 1954 and in Benghazi in 1960. These findings were based on questionnaires filled by middle age and elderly fishers (located in Tobrok, Ein Ghazala, At Timimi, Ras Ettin, Darna, Khalij Bumba, Susah, Benghazi fishing sites on eastern Libya), on reports and documents of “Darna Fishers Trade Union”, “Tobrok Fishers Trade Union” and “Office of Marine Resources in Darna” and on consultations with members of these institutions. Estimates of monthly catch from the two *Siganids* per fisher per fishing site shortly after they established themselves in the eastern coast ranged between 2 to 60 kg (average = 17.25 kg) with *S. luridus* been more abundant than *S. rivulatus*. Both fishes are residents of shallow coastal water, 1 – 8 meters deep (average = 4.25 meters). Average total length of the initial population the two *Siganids* was 23.44 cm. Questionnairing fishers, in particularly elderly ones, is a powerful new technique for establishing more accurate records of first entry than that given by scientist. However, this technique is limited by the scarcity in elderly fishermen who are inventory of information of the old times but are unfortunately eliminated through time. Yet the approach may not be too late for many Lessepsian fishes and scientists are encouraged to use it.

KEYWORDS: Lessepsian migrants, dusky spinefoot fish, *Siganus luridus*, marbled spinefoot, *Siganus rivulatus*, eastern Libya, Mediterranean Sea

INTRODUCTION

Both the Mediterranean Sea and the Red Sea are semi closed basins that are connected to adjacent oceans by narrow and shallow gates, the Mediterranean Sea to the Atlantic ocean

through the strait of Gibraltar, and the Red Sea to the Indian Ocean through the Gulf of Aden and Bab el Mandeb straits. In addition, the Red has acquired an artificial connection to the Mediterranean Sea via the Suez Canal since 1869. The Canal is 193km long, 24m deep and 205m wide. The 2014-2016 expansion and widening of the Canal doubled its ship passage capacity by allowing sailing in both directions.

The Canal provides a limited chance for Red Sea species to migrate to the Mediterranean Sea (Lessepsian migrants) and vice versa (anti-Lessepsian migrants). The number of Lessepsian migrant exceeds by far the number of anti-Lessepsian migrants (Cevik *et al.*, 2002). According to the Mediterranean Action Plan for Invasive Species UNEP-MAP-RAC/SPA, 2005; Bazairi *et al.*, 2013 and Katsanevakis *et al.*, 2014a,b, the main known pathway/vector of species introduction into the Mediterranean Sea is the Suez Canal followed by shipping (ballast water and sediments, anchoring and fouling), aquaculture (both marine and brackish species) and trade in live marine species (aquarium activities, fishing baits, seafood).

Golani, 2010, stated that frequently, invasive species are discovered after they have already been in the new habitat for extended period of time making it difficult to determine how the invasion started and when. The two Siganids rabbit fishes, the dusky spinefoot *S. luridus* (Ruppell, 1829) and the marbled spinefoot *S. rivulatus* (Forsskal, 1775) were recorded in Libya for the first time in 1968 and 1970 in order (Stirn, 1970; Lamboeuf, 2000; Bazairi *et al.*, 2013). However, some elderly Libyan fishers insisted that they have known the two fishes in the eastern coast since the beginning of the 1950ties. The objectives of the present study were, therefore, to clarify this contradiction by:

- i- Giving a comprehensive account on Lessepsian migration in the Mediterranean Sea with emphasis on Libya, in particular of the two Siganids: *S. luridus* and *S. rivulatus*.
- ii- Investigating date of first appearance of *S. luridus* and *S. rivulatus* in the eastern coast of Libya using:
 - questioners filled by some fishers of the eastern coast of Libya Mediterranean Sea.
 - reports and documents of “Darna Fishers Trade Union”, “Tobrok Fishers Trade Union” and “Office of Marine Resources in Darna” and on consultations with members of these institutions

LITERATURE REVIEW

Problems caused by aliens

In addition to alterations in ecosystem functioning and biodiversity loss, alien species can also cause serious economic (fisheries, tourism and aquaculture) and health impacts. Many of the marine alien species exhibit aggressive invasive behavior (Pimentel *et al.*, 2000; Vilà *et al.*, 2010; Bazairi *et al.*, 2013; Katsanevakis *et al.*, 2014a, b.). However it is generally assumed that

lessepsian migrants do not seem to influence the abundance of native species (Madl, 2011). This particularly worrying at today rate of introduction into the Mediterranean Sea of 1 new species every 10–14 days (Zenetos, 2010; Zenetos *et al.*, 2012; Galil *et al.*, 2014).

Mitigation measures for introducing alien species

Mitigation measures for introduction of aliens in the Mediterranean Sea were discussed by Galil, 2007. Although many international laws, conferences, agreements and treaties (e.g. CBD, 1992, Article 8h; IMO; UNCLOS, 1982; Barcelona Convention, 1995; SPA, 1982; UNEP; RAC/SPA) have called for or adopted such measures yet progress is slow and rudimentary and an effective policy of prevention is hardly enforced (Occhipinti-Ambrogi, 2004), even the appreciation of the problem by politicians and the causative sectors is limited.

Indigenous, alien and Lessepsian biota of the Mediterranean Sea

The biota of the Red Sea is of tropical Indo-Pacific origin, while that of the Mediterranean is primarily of Atlanto-Mediterranean species (62%) derived from the adjacent Atlantic biogeographic provinces beyond the strait of Gibraltar. Many Mediterranean species are endemic (28%) (Shakman, 1969; Fredj *et al.*, 1992) while others are cosmopolitan, Circum-tropical (13%) or Indo-Pacific (5%). Boudouresque *et al.*, 2005, reported flora and fauna of the Mediterranean Sea as around 12 000 species. Coll *et al.*, 2010, reported 17000. Based on Zenetos *et al.*, 2010, 2011a, b, 2012; Bazairi *et al.*, 2013, a total of 986 alien species were known in the Mediterranean by 2012: 775 in the Eastern basin, 308 in the Western basin, 249 in the Central basin and 190 in the Adriatic Sea. About 54% of these species are Lessepsian (Catsanevakis, 2012). According to Samuel and Frotté, 2015, the number of Mediterranean marine fishes is currently about 750 (Golani *et al.* 2013), almost 90 of them are Lessepsian migrant (Belmaker *et al.*, 2013; Golani *et al.*, 2002), with new species regularly being added to the list (Golani, 2006). Generally, the richness of Mediterranean species diversity including fishes decreases from the West to the East (Ketchum, 1983; Quignard and Tomasini, 2000). In contrast, there is a clear east to west gradient for Lessepsian species richness (Coll *et al.*, 2010). The majority of the farthest spread species are early settlers (Oral, 2010). An invasive species (= invasive alien) is an introduced species which is ecologically and/or economically harmful (Boudouresque and Verlaque, 2002a,b; Occhipinti-Ambrogi and Galil, 2004; Sala *et al.*, 2011; Giakoumi, 2014). On average, 10% of introduced species become invasive (Williamson and Fitter, 1996; Boudouresque and Verlaque, 2002a). However in certain areas of the eastern Mediterranean some alien fishes are commercially exploited (Kallianiotis and Lekkas, 2005; Giakoumi, 2014; Tsikliras and Stergiou, 2014).

Factors that limit/facilitate introduction and spread of Lessepsian migrants

These factors were discussed by Shakman, 1969; Ben-Tuvia, 1972; Golani, 1996; Golani, 1998; Kovalev, 2006; Ben Rais *et al.*, 2008; Madl, 2011.

1- Narrowness and shallowness of the canal

The Suez canal is not conducive to Lessepsian migration because it is shallow, narrow, very long, lack rocky bottom substrates and polluted from maritime activity (Por, 1978; Golani, 1998). However, it has been substantially deepened and broadened (Kovalev, 2006) during the 1960s, the 1980s and 2014-2016.

2- The hypersaline lakes

Just after completion of the Suez Canal in 1869, Lessepsian migration was blocked by the presence of hypersaline lakes (a remnant of the once landlocked evaporated seas that connected the Mediterranean to the Red Sea) inside the canal. After the salinity of these lakes equalized through years, the barrier is eliminated and species began migrating into the Mediterranean Sea (George and Athanassiou, 1967; Kovalev, 2006; Ben Rais *et al.*, 2008; Madl, 2011) via the Suez Canal.

3- Aswan Dam: Decreased salinity at Port Saeed

Before construction of Aswan dam across the Nile River in 1964, floodwater used to spread as a surface layer off the northern entrance of the Canal (off Port Saeed). This low salinity water acted as a barrier to lessepsian migrants. After the construction the dam influx from the Nile was reduced and the salinity barrier disappeared and Lessepsian migration is facilitated (Galil, 2006; Kovalev, 2006; Ben Rais *et al.*, 2008; Madl, 2011).

4- Prevailing temperature

Red sea organisms are tropical; the Mediterranean environment is subtropical-temperate and therefore is not hospitable to Lessepsian species. However, the global rise in temperatures during the present and the last century (IPCC, 2013; Jones *et al.*, 2013; Marras *et al.*, 2015) has led to “tropicalization” of the Mediterranean, resulting in the present day high rate of introduction of Lessepsian species (Bianchi, 2007; Ben-Tuvia, 1972; Raitos *et al.*, 2010; Madl, 2011; Hiddink *et al.*, 2012; Bazairi *et al.*, 2013; Marras *et al.*, 2015; Libralato *et al.*, 2015).

5- Salinity

Salinity is higher in the Red Sea than in the Mediterranean. This made it easier to Lessepsian migrants to flourish in the Mediterranean Sea (Ben-Tuvia, 1966; Kovalev, 2006).

6- Hydrostatic Pressure/currents

The general northwardly flow of water along the Suez Canal caused by prevailing winds and currents and the higher surface level of the Red Sea compared to that of the Mediterranean Sea has aided Lessepsian migration and attenuated anti-Lessepsian migration (Hassan *et al.*, 2003; Ben Rais *et al.*, 2008; Madl, 2011).

7- Oligotrophic

Red Sea fauna are used to living in nutrient poor environments of the Red Sea, and therefore, are able to flourish in the relatively nutrient rich oligotrophic waters of the Mediterranean.

8- Time factor

Immigration is a continuous process, and over time the probability of suitable species of fishes entering the Suez Canal and colonizing the new region increases (Ben-Tuvia, 1972).

9- unsaturated niche

The successful colonization of the Mediterranean by Lessepsian Migrants is in some cases facilitated by their ability to compete with local species on the same resources and by filling an unsaturated niche in their new habitat (Madl, 2011).

Potential migrants are “R-strategist” (Madl, 2011).

10- The east west gradient

The east west gradient of invasive species was studied by Golani 1998. Time is a factor in westward distribution. At present there is no conclusive explanation why some species continue westward and others do not. According to Harmelin-Vivien and Harmelin, 1990; Wootton, 1992 and the combination of temperature regime and thermal tolerance of fish species determines distribution.

11- The Gulf of Suez, the Suez canal and the south eastern Mediterranean Sea are temporary “resting” (acclimation) sites for lessepsian migrants on their way. From the latter site they can spread to other parts of the Mediterranean Sea. The Mediterranean Sea also has its own resting sites. Resting sites do not work well in the opposite direction (anti-Lessepsian migration).

Alien marine species of Libya

The list of alien species of Libya Mediterranean was updated by Bazairi *et al*, 2013 (but see Stirn, 1970; Zupanovic and El-Buni, 1982; Al-Hassan and El-Silini, 1999; Lamboeuf, 2000; Ben-Abdallah *et al*, 2005): until now 63 marine aliens species have been recorded along the Libyan coastline, the majority been of Indo-Pacific origin (53%), followed by Red Sea – Indian Ocean species (16%), Red Sea species (14%), Circum-tropical (3%) and less than 5% in the Atlantic (Bazairi *et al*. 2013).

The through studies of Lessepsian fish in Libya by Shakman and Kinzelbach (e.g. Shakman, 1969; Shakman and Kinzelbach, 2007) showed that sixteen Lessepsian fish species, representing 14 families, were found in Libya versus 107 in Tunisia and 232 in Egypt (Katsanevakis *et al.*, 2012; UNEP-MAP-RAC/SPA, 2013). The low number in Libya is probably an underestimate due to the limited effort of investigation (Shakman, 1969) the percentage of exotic fish species relative to native fish species decreases from east to west along the Libyan coast. Approximately 50% of the Libyan lessepsian migrants were found along the entire coast, 12.5% in the east and central regions, and 37.5% restricted to the eastern coast (Shakman, 1969). This means that there is a correlation between early arrival and greater abundance which can be explained (Shakman and Kinzelbach, 2007): (a) because the longer a species is in the Mediterranean, the greater the opportunity to build up its population, or (b) by the greater research effort, which was much less

intense in the past (Golani, 2002). More than 37% of the recorded Lessepsian fish species in Libya are of commercial value, especially rabbitfish (Shakman, 1969).

***S. luridus* and *S. rivulatus*: General description**

S. luridus and *S. rivulatus* are medium size coastal fishes. Body ellipsoid, oval and compressed. Their maximum lengths are about 30cm for the former and 27 cm for the latter but a more common length for both is 20 cm (Sommer *et al.*, 1996; Bariche, 2002). The meristic formulae are: D, XIV+10; A, VII + 8-9; P, 16-17; V, I+3+I and D, XIV+10; A, VII + 8-10; P, 15-17; V, I+3+I in order (Golani, 2006). Both fishes are often found in small schools feeding on benthic algae in shallow water with compacted sand substrates. The dorsal and anal fin spines contain venom that is not life-threatening to humans but causes severe pain (FishBase, WRMS).

Distribution of *S. luridus* and *S. rivulatus* in the Mediterranean Sea and in Libya

The two Siganids, *S. luridus* and *S. rivulatus* are widely distributed in the Indian Ocean and the Red Sea, respectively (Bazairi *et al.*, 2013). They were initially found in the Levantine basin, but later on they spread to the north and west where they became established in the central Mediterranean and constituted important fisheries (Stirn, 1970; Ktari and Ktari, 1974; Golani *et al.*, 2002; Ben Abdallah *et al.*, 2003; Streftaris and Zenetos, 2006; Shakman *et al.*, 2008). Their spread from the east to different parts of the Mediterranean sea was reviewed by Shakman, 1969; Ismen *et al.*, 2015. The two fishes were first recorded in Palestine in 1927 (Steinitz, 1927), then off the coast of Syria in 1931 (Gruvel, 1931). *S. luridus* was recorded in Libya in 1968 (Stirn, 1970; Lamboeuf, 2000; Bazairi *et al.*, 2013), in Tunisia in 1971, but crossed the Strait of Sicily only in 2004; it reached the French coast in 2008 (Ktari and Bouhalal, 1971; Ktari and Ktari, 1974; Daniel *et al.*, 2009). The most recent record was in the Aegean Sea in 2014 (Ismen *et al.*, 2015). The first record of *S. rivulatus* in Libya was in 1970 (Stirn, 1970).

In Libya *S. luridus* seems to be more present in central and western parts of the littoral (Shakman and Kinzelbach, 2007c) while *S. rivulatus* is mostly common in the east. Both fishes are common in the artisanal catch of Libya but their abundance decrease towards the west. They are captured by trammel and gill nets (Ben Abdallah *et al.*, 2003; Galil, 2006; Bazairi *et al.*, 2013; Khamis, 2008). The rapid westward spread of the two Siganids can be explained (Bariche *et al.*, 2003; Castriota and Andaloro, 2005; Bianchi, 2007; Corsini-Foka *et al.*, 2010; Poloniato *et al.*, 2010; Por, 2010; Raitsos *et al.*, 2010; Marras *et al.*, 2015):

- The rise in water temperature of the whole Mediterranean made this sea more receptive to Lessepsian emigrants which are of tropical origin.
- Introduced Siganids, for example *S. luridus*, successfully outcompetes its main indigenous counterpart herbivores (Bariche *et al.*, 2004)
- The increased productivity of the Mediterranean as one moves from east towards west favors the westward spread of Lessepsian migrants, especially herbivores.

- Eco-physiological plasticity: Its spawning season is shortened, probably in relation to the varied seawater temperatures that are found in the Mediterranean Sea (Bariche *et al.*, 2003; Castriota and Andaloro, 2005).

METHODS

First appearance of *S. luridus* and *S. rivulatus* in the artisanal catch from the eastern coast of Libya Mediterranean Sea was investigated by two methods:

i- By questionnaires filled by 50 fishers

These fishers were located in the major fish landing sites of the eastern coast of Libya. The study area (Figure 1) extends from Burde in the east of Libya coast to Breka in the west, an approximately 500km shore line. The landing sites investigated were Tobrok, Ein Ghazala, At Timimi, Ras Ettin, Darna, Khalij Bumba, Susah, Benghazi. The questionnaire was filled by six to seven fishers present in each of these sites. Most fishers questionnaired were 60 years old and above. Names, locations, addresses and cell phone numbers of the chosen fishers were provided by “Darna Fishers Trade Union”, “Tobrok Fishers Trade Union” and “Office of Marine Resources in Darna”. These institutions also helped in delivering the questionnaires to individual fishers and in retrieving them after been filled. The questionnaire contained the following questions:

- What is the name of the area you fish in or used to fish in?
- When did you first see *S. luridus* and *S. rivulatus* in your catch?
- What gear is most effective in catching *S. luridus* and *S. rivulatus*?
- At what depth is *S. luridus* and *S. rivulatus* are most common?
- Give average length and weight of *S. luridus* and *S. rivulatus*.
- Rate abundance of *S. luridus* and *S. rivulatus* in your catch: (-) rare, (1) present, (2) common, (3) abundant.

The fishers, however, stipulated that we do not mention any of their personal information including names and ages in our study.




Figure 1. Map of the shore line of the eastern coast of Libya (the study area)

- ii- **By investigating reports and records available in old files of** “Office of Marine Resources in Darna”, “Darna Fishers Trade Union” and “Tobrok Fishers Trade Union” for first mentioning of *S. luridus* and *S. rivulatus*, and on consultations with members of these institutions. The activities of these three institutions covers all the eastern coast of Libya (the study area).

RESULTS

Records of first appearance of *S. luridus* and *S. rivulatus* in the artisanal catch of the eastern coast of Libya is documented in Tables 1 to 5. Data presented in these tables were compiled by integrating data obtained from the questionnaire with that obtained from the “Office of Marine Resources in Darna” and the Fishers Trade Unions of Darna and Tobrok. The two Siganids were first observed in Libya in Tobrok in 1950, in Darna in 1954 and in Benghazi in 1960 (Table 1). These three sites are the most prominent fishing and landing sites of the eastern coast of Libya. The two new comers appeared simultaneously in the catch for the first time. Elderly fishers from Tobrok, Darna and Benghazi still recall their painful stings when they attempted to remove them

from their nets and lines, the main fishing gears in use at that date. Initially they were called “Alsamakah um Shoak” meaning in Arabic “the fish with spines” and were not marketed but thrown away. However, after their numbers increased in the catch they became acceptable by local consumers; *S. luridus* soon became known locally as “Batata soada” (Arabic for “black potatoes”) and *S. rivulatus* as “Batata baida” (“white potatoes”).

Table 1. Records of first appearance of *S. luridus* and *S. rivulatus* in the studied sites.

location	Geographic coordinates		Fishing gear	Record date
	Latitude	Longitude		
Tobrok	N 23°55'	E '07°32	Gill net	1950 – 1960
Ein Ghazala	N 23°55'	E '07°32	small hooks, Gill net	1967-1975
At Timimi	N 23°06'	E '19°32	Hand line, gill net	1970 -1980
Ras Ettin	N 23°07'	E '37°32	Gill net	1960 -1980
Darna	N 22°39'	E '16° 32	Gill net	1954 - 1960
Khalij Bumba	N 23°06'	E '25°32	small hooks	1990
Susah	N 21° 58'	E '54°31	Gill net	1970
Benghazi	N 20°03'	E '36°32	Gill net	1960 - 1965

Estimates of monthly catch from the two Siganids per fisher per fishing site (Table 2 and Figure 2) shortly after they established themselves in the eastern coast ranged between 2 to 60 kg (average = 17.25 kg) with *S. luridus* been more abundant than *S. rivulatus* (Table 3 and Figure 3). Both fishes are residents of shallow coastal water (Table 4 and Figure 4), 1 to 8 meters deep (average = 4.25 meters).

Table 2. Estimates of monthly catch from *S. rivulatus* and *S. luridus* per fisher per site shortly after their introduction.

Fishing/Landing site	Monthly catch in kg per fisher from <i>S. luridus</i> and <i>S. rivulatus</i>	
	Average	Range
Tobrok	7.5	5 - 10
Ein Ghazala	6.5	3 - 10
At Timimi	22	4 - 40
Ras Ettin	7.5	6 - 9
Darna	32.5	5 - 60
Khalij Bumba	16.5	3 - 30
Susah	13.5	2 - 25
Benghazi	32	4 - 60
Grand average catch in kg/month/fisher/site		17.25




Figure 2. Estimates of monthly catch from *S. rivulatus* and *S. luridus* per fisher per site shortly after their introduction.

Table 3. Relative abundance of *S. luridus* and *S. rivulatus* in the studied sites. (-) rare, (1) present, (2) common, (3) abundant

Fishing/Landing site	Relative abundance	
	<i>S. luridus</i>	<i>S. rivulatus</i>
Tobrok	2	1
Ein Ghazala	2	2
At Timimi	3	2
Ras Ettin	3	2
Darna	2	-
Khalij Bumba	2	1
Susah	3	-
Benghazi	1	2
Grand average length	3.35	1.25





Figure 3. Relative abundance of *S. luridus* and *S. rivulatus* in the studied sites.

Table 4. Depth of occurrence of *S. luridus* and *S. rivulatus* in the studied sites.

Fishing/Landing site	Depth of <i>S. luridus</i> and <i>S. rivulatus</i> in m	
	Average	Range
Tobrok	4	3 - 5
Ein Ghazala	4	1 - 7
At Timimi	4.5	1 - 8
Ras Ettin	6	4 - 8
Darna	2.5	2 - 3
Khalij Bumba	3	2 - 4
Susah	4.5	4 - 5
Benghazi	5.5	4 - 7

Figure 4. Depth of occurrence of *S. luridus* and *S. rivulatus* in the studied sites.

Fishers did not give separate fish lengths for *S. luridus* and *S. rivulatus* when first observed in the catch but gave a common average estimates for both fishes (Table 5 and Figure 5). Common lengths of *S. luridus* and *S. rivulatus* ranged between 10 to 40 cm with an average of 23.44 cm.

Table 5. Average total lengths of *S. luridus* and *S. rivulatus* in the studied sites. The fishers gave a common length estimate for both fishes. They were not able to give separate estimates for each fish.

Fishing/Landing site	Total lengths of <i>S. luridus</i> and <i>S. rivulatus</i> in cm	
	Average	Range
Tobrok	25	15 - 35
Ein Ghazala	20	10 - 30
At Timimi	25	15 - 35
Ras Ettin	25	20 - 30
Darna	22.5	15 - 30
Khalij Bumba	25	20 - 30
Susah	25	15 - 35
Benghazi	20	10 - 30
Grand average length	23.44	




Figure 5. Average total lengths of *S. luridus* and *S. rivulatus* in the studied sites. The fishers gave a common length estimate for both fishes.

DISCUSSION

Establishing dates of first introduction of lessepsian migrants into different parts of Mediterranean Sea is important for studying biology, ecological relationships and acquired genetic variations of the organisms in their new habitat (Giacomo, 2014). Golani, 2010, stated that frequently, invasive species are discovered after they have already been in the new habitat for extended period of time making it difficult to determine how the invasion started and when.

According to Stirn, 1970; Lamboeuf, 2000; and Bazairi *et al.*, 2013 *S. luridus* and *S. rivulatus* were recorded in Libya for the first time in 1968 and 1970 in order. However, the present study shows that the two Siganids invaded Libya at a much earlier date: in Tobrok in 1950 and in Darna in 19954. Questionnaire is a powerful technique for establishing date of first introduction. However, this technique is limited by the scarcity in elder fishermen who are inventory of information of the old times but are unfortunately eliminated through time. This may explain the much more recent dates of introduction into Ein Ghazala (1967), At Timimi (1970), Ras Ettin (1960), Khalij Bumba (1990), Susah (1970) and Benghazi (1960).

From the present study it appears that the two Siganids can grow into sizable populations and contribute to the catch shortly after they introduce themselves into a new site, with *S. luridus* becoming the more abundant one; more than twice abundant than *S. rivulatus*. This is in agreement with Shakman *et al.*, 2008 who mentioned that the two species are abundant in the rocky shallow Libyan coasts and contribute significantly to the catch. One reason for the

prosperity of these two herbivores is the richness of the Libyan coast in seaweeds and sea grass; another reason is their ability to compete with local herbivores.

Both Siganids inhabit shallow coastal waters: 1- 8 meters (average = 4.25 meters) where they are caught by nets and lines by fishers using boats or wadding.

According to the questionnaired fishers the total length of the two species at the time they were practicing fishing ranged between 10 to 35 cm, with an average of 23.44 cm. Thirty five centimeters is more than the length recorded presently for the two Siganids in the international literature. Sommer *et al.*, 1996; Bariche, 2002, 2005, reported that maximum lengths were 28 cm for *S. luridus* and 30 cm for *S. rivulatus* from the Mediterranean Sea and the Red Sea but a more common length for both is 20 cm.

The implication of the present study to scientific approach is that the date of first entry to a particular region attributed by a scientist(s) to a Lessepsian fish may be lagging decades behind the true date of entry. When a scientist discovers a new invasive fish he/she must ask the local fishers, in particular the elderly ones, when they have first become acquainted with the fish. This is may not be too late for many "foreign" fishes in many regions.

REFERENCES

- Al-Hassan L.A.J., El-Silini O. A., (1999). Check-list of bony fishes collected from the Mediterranean coast of Benghazi, Libya. *Revista de Biologia Marina y Oceanografia* 34: 291–301.
- Bariche, M., (2002). Biologie et écologie de deux espèces lessepsiennes (*Siganus rivulatus* et *Siganus luridus*, Téléostéens, Siganidae) sur les côtes du Liban. Thèse de Doctorat, Université de la Méditerranée (Aix-Marseille II). 223 p
- Bariche, M. (2005). Age and growth of Lessepsian rabbitfish from the eastern Mediterranean. *Journal of Applied Ichthyology* (Impact Factor: 0.87). 04/2005; 21:141-145. DOI: 10.1111/j.1439-0426.2004.00619.x
- Bariche M, MH Vivien and J. P. Quignard, (2003). Reproductive cycles and spawning periods of two Lessepsian Siganid fishes on the Lebanese coast. *Journal of Fish Biology* 62: 129-142.
- Bariche M, Y Letourneur and M. H. Vivien, (2004). Temporal fluctuations and settlement patterns of native and Lessepsian herbivorous fishes on the Lebanese coast (eastern Mediterranean). *Environmental Biology of Fishes* 70: 81-90.
- BAZAIRI H., Y. R. SGHAIER, I. BENAMER, H. LANGAR, G. PERGENT, E.M. BOURASS, M. VERLAQUE, J. BEN SOUSSI and A. ZENETOS, (2013). Alien marine species of Libya: first inventory and new records in El-Kouf National Park (Cyrenaica) and the neighboring areas. *Medit. Mar. Sci.*, 14/2, 2013, 451-462 451
- Belmaker J, Parravicini V, Kulbicki M (2013) Ecological traits and environmental affinity explain Red Sea fish introduction into the Mediterranean. *Glob Change Biol* 19: 1373–1382.

- Ben Abdallah, A. R., Al-Turky A., Fituri, A. A., (2003). Commercially exploited exotic fishes in Libya. Bulletin de l'Institut National des Sciences et Technologies de la Mer (Salammbô) n°spécial, (8), Actes des 6èmes Journées de l'ATSMer, Tunis (Tunisie) 28-30 Novembre 2003, 53-55.
- Ben Abdallah, A. R., Al-Turky A., Fituri, A. A., (2005). Exotic fishes along the coast of Libya (Arabic with English abstract) Libyan Journal of Marine Science, (10), 48-56.
- Ben Rais Lasram F., J. A. Tomasini, F. Guilhaumon, M. S. Romdhane, T. Do Chi, D. Mouillot, (2008). Ecological correlates of dispersal success of Lessepsian fishes. Mar Ecol Prog Ser Vol. 363: 273–286, 2008 doi: 10.3354/meps07474.
- Ben-Tuvia A., (1966). Red sea fishes recently found in the Mediterranean. Copeia 2: 254–275.
- Ben-Tuvia, A., (1972). Immigration of fishes through the Suez Canal. 17e Congr. Intern. Zool. No.3. Les conséquences biologiques des canaux inter-océans.
- Bianchi, C. N., (2007). Biodiversity issues for the forthcoming tropical Mediterranean Sea. Hydrobiologia, 580, 7-21.
- Boudouresque, C.F. and Verlaque, M., (2002a). Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. Mar. Poll. Bull., 44: 32-38.
- Boudouresque, C.F. and Verlaque, M. (2002b). Assessing scale and impact of ship-transported alien macrophytes in the Mediterranean Sea. In: Alien marine organisms introduced by ships in the Mediterranean and Black seas. Briand F. (ed.), CIESM Workshop Monographs, 20: 53-61.
- Boudouresque C. F., Sandrine Ruitton and Marc Verlaque, (2005). LARGE-SCALE DISTURBANCES, REGIME SHIFT AND RECOVERY IN LITTORAL SYSTEMS SUBJECT TO BIOLOGICAL INVASIONS. In Velikova v., Chipev N. eds Unesco-Roste BAS / Workshop on regime shifts, 14-16 June, Varna, Bulgaria, 85-101.
- Castriota I. and F. Andaloro, (2005). First record of the lessepsian fish *Siganus luridus* (Osteichthyes: Siganidae) in the Tyrrhenian Sea. Marine Biodiversity Records, page 1 of 2. #2005 Marine Biological Association of the United Kingdom doi:10.1017/S1755267205001223; Vol. 1; e11; 2008 Published online
- CEVIK C., TOKLU B., SARIHAN E., (2002) The impact of the Red sea species (Lessepsian) on the Mediterranean Sea. In: Bayram Öztürk and Nuri Basusta (eds). Workshop on Lessepsian migration, 20-21 July 2002 Gökceada, Turkey, 83-86.
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F, *et al.*, (2010) The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats. PLoS ONE 5(8): e11842. doi:10.1371/journal.pone.0011842
- Corsini-Foka M., Pancucci-Papadopoulou M.-A. and Kalogirou S., (2010) Is the Lessepsian province in expansion? The Aegean Sea experience. FAO-EastMed Working Document. Sub-regional technical meeting on the Lessepsian migration and its impact on eastern mediterranean fishery. Nicosia, 7–9 December 2010, pp. 50–59.
- Daniel B., Piro S., Charbonnel E., Francour P. and Letourneur Y., (2009) Lessepsian rabbitfish *Siganus luridus* reached the French Mediterranean coasts. Cybium 33, 163–164. Fishbase. Retrieved 15 June 2013.

- Forsskål, P., (1775). *Descriptiones animalium, avium, amphibiorum, piscium, insectorum, vermium; quae in itinere orientali observavit Petrus Forsskål. Post mortem auctoris edidit Carsten Niebuhr. Havenai, Mölleri.*
- FREDJ G., MEINARDI M., MENARDI M. (1992) État des Connaissances sur la Faune marine Méditerranéenne. Bulletin de l'Institut Océanographique de Monaco, no. Spécial 9: 133-145.
- GALIL B.S., (2006). The marine caravan – the Suez Canal and the Erythre: an invasion. In: Gollasch, B. Galil, A Cohen. *Bridging divides: maritime canals as invasion corridors.* Springer (Monographie Biologique, 83.). Part 4: The Suez Canal. pp. 207-300.
- Galil, Bella S. • Ferdinando Boero • Marnie L. Campbell • James T. Carlton • Elizabeth Cook • Simonetta Frascchetti • Stephan Gollasch • Chad L. Hewitt • Anders Jelmert • Enrique Macpherson • Agnese Marchini • Cynthia McKenzie • Dan Minchin • Anna Occhipinti Ambrogi • Henn Ojaveer • Sergej Olenin • Stefano Piraino • Gregory M. Ruiz., (2014). 'Double trouble': the expansion of the Suez Canal and marine bioinvasions in the Mediterranean Sea. *Biol Invasions* DOI 10.1007/s10530-014-0778-y
- George C. J., Athanassiou V., (1967) A two year study of the fishes appearing in the seine fishery of St George Bay, Lebanon. *Ann Mus Civ Stor Nat Genova* 76:237–294
- Giakoumi S., (2014) Distribution patterns of the invasive herbivore *Siganus luridus* (Ruppell, 1829) and its relation to native benthic communities in the central Aegean Sea, Northeastern Mediterranean. *Marine Ecology* 35, 96–105.
- Golani D., (1996). The marine ichthyofauna of the eastern Levant—history, inventory and characterization. *Isr. J. Zool.*, 42. 15-55.
- Golani, D., (1998). Distribution of Lessepsian migrant fish in the Mediterranean, *Italian Journal of Zoology*, 65:S1, 95-99, DOI: 10.1080/11250009809386801
- Golani D., (2002). Lessepsian fish migration—Characterization and impact on the Eastern Mediterranean. Pp. 1–9. In: Basusta N. (eds.) *Workshop on Lessepsian migration*, 20–21 July 2002, Groceada, Turkey.
- Golani D., (2006). The Indian scad (*Decapterus russelli*), (Osteichthyes: Carangidae), a new Indo-Pacific fish invader of the eastern Mediterranean. *Scientia Marina* 70: 603–605.
- Golani D., (2010). Colonization of the Mediterranean by Red Sea fishes via the Suez Canal Lessepsian Migration. Pp. 145–188. In: Golani D., Appelbaum-Golani B. (eds.) *Fish invasions of the Mediterranean Change and renewal.* Pensoft, Sofia, Bulgaria.
- Golani D., Orsi-Relini L., Massuti E. and Quignard J. P., (2002) CIESM atlas of exotic species in the Mediterranean. In Briand F. (ed.) *Fishes*, Vol. 1. Monaco: CIESM Publications, 259 pp.
- Golani D., Bayram Öztürk, Nuri Basusta, (2006). *Fishes of the Eastern Mediterranean.* Printed by: Graphis Matbaa +902126290607
- Golani D, Orsi-Relini L, Massuti E, Quignard JP, Dulčić J, Azzurro E (2013) CIESM atlas of exotic fishes in the Mediterranean. <http://www.ciesm.org/atlas/appendix1.html> (Accessed March 2015) Quignard JP, Tomasini JA (2000) Mediterranean fish biodiversity. *Biologia Marina Mediterranea* 7(3): 1–66

- Gruvel A., (1931) Les Etats De Syrie. Richness Marines et Fluviales. Exploitation Actuelle, Avenir. Paris: Societe d'editions Geographiques, Marines et Coloniales, pp. 72–134.
- Harmelin-Vivien M., Harmelin J. G., (1990) -Guide des Poissons dela Méditerranée. Delachaux et Niestlé, Lausanne, Switzerland, 144pp.
- Hassan, M., Harmelin-Vivien, M. & Bonhomme, F., (2003). Lessepsian invasion without bottleneck: example of two rabbitfish species (*Siganus rivulatus* and *Siganus luridus*). Journal of the Experimental Marine Biology and Ecology, 291, 219-232.
- Hiddink J. G., Ben Rais Lasram F, Cantrill J, Davies A. J. (2012). Keeping pace with climate change: what can we learn from the spread of Lessepsian migrants? Glob Change Biol18: 2161–2171.
- IPCC, (2013). Climate Change 2013: the Physical Science Basis. Working group 1 contribution to the fifth assessment report of the International Panel on Climate Change, International Panel on Climate Change, Cambridge, UK and New York, NY, USA.
- Ismen A., A. Ayaz and Z.D. Yildirim, (2015). Northernmost record of the dusky spinefoot *Siganus luridus* in the Aegean Sea (Turkey coast). Marine Biodiversity Records, page 1 of 4. Marine Biological Association of the United Kingdom, 2015. doi:10.1017/S1755267215000172; Vol. 8; e42; 2015 Published online.
- Jones, M. C., Dye, S. R., Pinnegar, J. K., Warren, R., and Cheung, W. W. L., (2013). Applying distribution model projections for an uncertain future: the case of the Pacific oyster in UK waters. Aquat. Conserv. Mar. Freshw. Ecosyst. 23, 710–722. doi: 10.1002/aqc.2364.
- Kallianiotis, A., and Lekkas, V., (2005). First documented report on the Lessepsian migrant *Etrumeus teres* De Kay, 1842 (Pisces: Clupeidae) in the Greek Seas. J. Biol. Res. 4, 225–229.
- Katsanevakis, S., Bogucarskis, K., Gatto, F., Vandekerckhove, J., Deriu, I. *et al.*, (2012). Building the European Alien Species Information Network (EASIN): a novel approach for the exploration of distributed alien species data. BioInvasions Records, 1, 235-245.
- Katsanevakis S, Coll M, Piroddi C, Steenbeek J, Ben Rais Lasram F, Zenetos A, Cardoso AC, (2014a). Invading the Mediterranean Sea: biodiversity patterns shaped by human activities. Frontiers in Marine Science, <http://dx.doi.org/10.3389/fmars.2014.00032>
- Katsanevakis S, Wallentinus I, Zenetos A, Leppäkoski E, Çinar ME, Oztürk B, Grabowski M, Golani D, Cardoso AC, (2014b). Impacts of marine invasive alien species on ecosystem services and biodiversity: a pan-European review. Aquatic Invasions9(4): 391–423.
- KETCHUM B.H., (1983). Ecosystems of the world 26 Estuaries and Enclosed Seas. Amsterdam, Oxford, New York: Elsevier Scientific Publishing Company.
- Khamis, E., (2008). The analysis of the catch trawling net along Ben Gghazi, M. sc, Thesis. CoastFac. of. Nat. Res. and Envi, Omar El Mukhtar University PP89.
- KOVALEV, (2006). On the problem of Lessepsian migrations of zooplanktonic organisms. Medit. Mar. Sci, 7/2, 2006, 67-71
- Ktari-Chakroun F. and Bouhalal M., (1971). Record of *Siganus luridus* (Ruppel) in the Gulf of Tunisia. Bull. Inst. Natl. Sci. Tech. Oceanogr. Peche Salammbô2, 49–52.

- Ktari F., Ktari M. H., (1974). Presence dans le Golf de Gabes de *Siganus luridus* (Rüppell, 1829) et de *Siganus rivulatus* (Forsskal, 1775) (Poissons, Siganidae) parasité par *Pseudohaliotrema polymorphus*. Bulletin de l'Institut national scientifique et technique d'Océanographie et de pêche de Salammb 3: 95–98.
- Lamboeuf M., (2000). Artisanal fisheries in Libya – census of fishing vessels and inventory of artisanal fishery métiers. Final Report, FAO- COPEMED-MBRC , 42 pp.
- Libralato, S., Alberto Caccin and Fabio Pranovi. (2015). Modeling species invasions using thermal and trophic niche dynamics under climate change. Front. Mar. Sci., 27 May 2015 | <http://dx.doi.org/10.3389/fmars.2015.00029>
- Madl P., (2011). Colloquial Meeting of Marine Biology I Headed by: dr. A. Goldschmid. Essay about the phenomenon of Lessepsian Migration.
- Marras, S., Andrea Cucco, Fabio Antognarelli, Ernesto Azzurro, Marco Milazzo, Michel Bariche, Momme Butenschön, Susan Kay, Massimiliano Di Bitetto, Giovanni Quattrocchi, Matteo Sinerchia and Paolo Domenici, (2015). Predicting future thermal habitat suitability of competing native and invasive fish species: from metabolic scope to oceanographic modelling. Conservation physiology. Volume 3 2015.
- Occhipinti-Ambrogi, A. and Galil, B. S., (2004). A uniform terminology on bioinvasions: a chimera or an operative tool? Mar. Poll. Bull., 49: 688-694.
- ORAL, M., (2010). Alien fish species in the Mediterranean – Black Sea Basin. J. Black Sea/Mediterranean Environment Vol. 16(1): 87-132 (2010)
- Pimentel, D., Lach, L., Zuniga, R., Morrison, D., (2000). Environmental and economic costs of non-indigenous species in the United States. BioScience, 50, 53-65.
- Poloniato, D., S. Ciriaco, R. Odorico, J. Dulcic, L. Lipej, (2010). First record of the dusky spinefoot *Siganus luridus* (Rüppell, 1828) in the Adriatic Sea. ANNALES . Ser. Hist. nat . 20 . 2010 . 2.
- Por F. D., (1978). Lessepsian migration - The influx of Red Sea biota into the Mediterranean by way of the Suez Canal. Ecological Studies, Vol. 23. Berlin-Heidelberg-New York: Springer-Verlag, pp. 88–194.
- Por F. D., (2010). The new Tethyan ichthyofauna of the Mediterranean- historical background and prospect. In Golani D. and Appelbaum-Golani B. (eds) Fish invasions of the Mediterranean Sea: change and renewal. Sofia-Moscow: Pensoft Publishers, pp. 13–33.
- QUIGNARD J. P., TOMASINI J. A., (2000). Mediterranean fish biodiversity. Biologia Marina Mediterranea 7: 1-66.
- Raitsos D. E., Beaugrand G., Georgopoulos D., Zenetos A., Pancucci-Papadopoulou A. M., Theocharis A. and Papathanassiou E., (2010). Global climate change amplifies the entry of tropical species into the Eastern Mediterranean Sea. Limnology and Oceanography 55, 1478–1484
- Rüppell, W. P. E. S., (1829). Atlas zu der Reise im nördlichen Afrika. Zoologie; Fische des rothen Meeres. (1828-1830).

- Sala, E., Kizilkaya, Z., Yildirim, D., Ballesteros, E., (2011). Alien marine fishes deplete algal biomass in the Eastern Mediterranean. *PLoS ONE* 6 (2), e17356. doi:10.1371/journal.pone.0017356.
- Samuel P. Iglésias and Lou Frotté, (2015). Alien marine fishes in Cyprus: update and new records. *Aquatic Invasions* (2015) Volume 10, Issue 4: 425–438 doi: <http://dx.doi.org/10.3391/ai.2015.10.4.06>
- SHAKMAN, (1969). Lessepsian Migrant fish Species of the Coastal Waters of Libya: Status, Biology, Ecology. MSC. Thesis. Universitat Rostock. Germany.
- Shakman E. A., (2008). Lessepsian Migrant fish species of the coastal waters of Libya: status, biology, ecology. Rostock, Germany: Institut Biowissenschaften Biodiversitätsforschung Allgemeine & Spezielle Zoologie, Rostock University, 130 pp.
- Shakman E. A. and R. Kinzelbach, (2007a). Distribution and characterization of Lessepsian migrant fish along the coast of Libya. *Acta Ichthyologica et Piscatoria* 37(1): 7-15.
- Shakman, E., and Kinzelbach, R., (2007b). Commercial fishery and fish species composition in coastal waters of Libya. *Rostocker Meeresbiologische Beiträge*, 18, S. 63-78.
- Shakman, E., Winkler, H.M., Oeberst, R., Kinzelbach, R., (2008). Morphometry, age and growth of *Siganus luridus* Rüppell, 1828 and *Siganus rivulatus* Forsskal, 1775 (Siganidae) in the central Mediterranean (Libyan coast). *Revista de Biologia Marina & Oceanografia*, 43 (3), 521-529
- Sommer, C., W. Schneider and J.-M. Poutiers, (1996). FAO species identification field guide for fishery purposes. The living marine resources of Somalia. FAO, Rome. 376 p.
- Steinitz, W., (1927). Beiträge zur Kenntnis der Küstenfauna Palastinas. I. Pubbl. Staz. Zool. Napoli 8, 331–353.
- Stelios Catsanevakis, (2013). Invading the Mediterranean Sea: pathways of introduction, distribution patterns and impact of alien species. In: MARBIGEN Conference on Mediterranean Marine Biodiversity in view of in view of climate change and invasion by alien species. 7-9 October 2013., Heraklion, Greece.
- Štirn, J., (1970). Some notes on western trends of Lessepsian migration. In: Journées Ichthyologiques, Rome, 30 Nov- 1 Dec., pp. 187-190, CIESM, Monaco.
- Streftaris, N., Zenetos, A., (2006). Alien marine species in the Mediterranean - the 100 ‘worst invasives’ and their impact. *Mediterranean Marine Science*, 7 (1), 87-118.
- Tsikliras, A. C., and Stergiou, K. I., (2014). The mean temperature of the catch increases quickly in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 515, 281–284. doi: 10.3354/meps11005
- UNEP-MAP-RAC/SPA, (2005). Action Plan concerning species introductions and invasive species in the Mediterranean Sea . RAC/SPA (Ed.), Tunis, 30 pp.
- UNEP- MAP-RAC/SPA - Tunis, (2013). United Nations. Environment 012013/06/ - Nations Environment Programme / Mediterranean Action Plan (UNEP/MAP).
- Vilà M, Basnou C, Pyšek P, Josefsson M, Genovesi P, Gollasch S, Nentwig W, Olenin S, Roques A, Roy D, Hulme PE, (2010). How well do we understand the impacts of alien species on ecosystem services? A pan-European, cross-taxa assessment. *Frontiers in Ecology and the Environment* 8(3): 135–144.

- Williamson, M., Fitter, A., (1996). The varying success of invaders. *Ecology*, 77(6): 1661-1666.
- Wootton R. J., (1992). *Fish ecology*. Blackie & Son Ltd., Glasgow, 212pp. World Register of Marine Species. Retrieved 15 June 2013
- Zenetos A., (2010). Trend in aliens species in the Mediterranean. An answer to Galil, 2009 «Taking stock: inventory of alien species in the Mediterranean Sea». *Biol. Invasions*, sous presse.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M. E., Garcia Raso, J. E. *et al.*, (2010). Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2), 318-493.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J. E. *et al.*, (2011a). Errata to the review article (*Medit. Mar. Sci.*, 11 (2): 318-493). Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 12 (2), 509-514.
- Zenetos A., Katsanevakis S., Poursanidis D., Crocetta F., Damalas D., Apostolopoulos G, Gravili C., Vardala-Theodorou E. and Malaquias M., (2011b). Marine alien species in Greek Seas: additions and amendments by 2010. *Mediterranean Marine Science* 12, 95–120. Correspondence should be addressed to: A. Ismen Faculty of Marine Science and Technology Canakkale Onsekiz Mart University, TR-17100, Turkey.
- Zenetos A, Gofas S, Morri C , Rosso A, Violanti D, García Raso JE, Çinar ME, Almogi-Labin A, Ates AS, Azzurro E, Ballesteros E, Bianchi C. N., Bilecenoglu M, Gambi MC, Giangrande A, Gravili C, Hyams-Kaphzan O, Karachle PK, Katsanevakis S, Lipej L, Mastrototaro F, Mineur F, Pancucci-Papadopoulou M. A., Ramos Esplá A, Salas C, San Martín G, Sfriso A, Streftaris N, Verlaque M., (2012). Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science* 13(2): 328–352, <http://dx.doi.org/10.12681/mms.327>
- Zupanovic S., El-Buni A. A., (1982). A contribution to demersal fish studies off the Libyan coast. *Bulletin of Marine Research Centre (Tripoli-Libya)* 3: 77–122.

Acknowledgements

Thanks are expressed to “Darna Fishers Trade Union”, “Tobrok Fishers Trade Union” and “Office of Marine Resources in Darna” for helping in filling the questionnaires by fishers, availing to us relevant documents and records of their institutions and the productive discussions and suggestions they made.