

PROXIMATE, MINERAL AND FUNCTIONAL PROPERTIES OF DEFATTED AND UNDEFATTED CASHEW (*ANACARDIUM OCCIDENTALE LINN.*) KERNEL FLOUR.**Emelike N.J.T, Barber L.I and Ebere C.O**

Department of Food Science and Technology, Rivers State University of Science and Technology, Port Harcourt, Rivers State, Port Harcourt, Nigeria.

ABSTRACT: *Defatted and undefatted cashew (Anacardium occidentale Linn.) kernel flour was produced to analyse their effect on the proximate, mineral and functional properties. It was ascertained that the proximate content of protein, crude fibre and carbohydrate (34.0, 6.2 and 32.2%, respectively) increased significantly in the defatted compared to undefatted cashew kernel flour. A decrease was observed in the moisture content, ether extract (crude fat) and total ash (4.4, 1.6 and 1.8%, respectively) of the defatted than the undefatted flour. There was a significant difference ($p < 0.05$) in all the analysed mineral elements of defatted to the undefatted flours except manganese with significantly higher ($p < 0.05$) value in the undefatted compared to the defatted sample (9.0 and 2.9mg/100g), respectively. There was no significant difference ($p > 0.05$) in the bulk density, foam capacity/stability between 30 min – 1h, emulsion capacity and nitrogen solubility at pH₉ of the defatted and undefatted cashew kernel flour. A significant difference was observed between these two samples in the water/fat absorption capacity, emulsion stability from 30 min – 1h and nitrogen solubility at pH₈. The results obtained in this study indicates that defatted cashew kernel flour is a good source of protein and can be substituted to wheat flour in snack production and formulation of children's meals to reduce the importation of wheat flour into Nigeria, can be a useful source of low-fat fabricated foods as well as animal feeds as the ash content meets up the recommended allowance for animal feeding.*

KEYWORDS: Defatted/Undefatted Cashew Kernel Flour, Quality Characteristics

INTRODUCTION

Current trend in nutrition is the consumption of functional foods “foods that not only supply basic nutrients but also help to prevent certain diseases” advocated by World Nutrition Bodies due to different health related problems with wheat flour consumption such as celiac disease, diabetes and coronary heart disease (WHO/FAO, 2003). This situation has created the need for the consumption of low-carbohydrate diets, slowly digested starchy foods as well as an increased intake of functional foods (Hurs and Martins, 2005). Many plants usually in the form of protein extracts, fibre extract or seed flours are being investigated and tested for new products such as low cost fabricated foods with high nutritional value, attractive and acceptable to consumers. Snack products such as biscuits and cookies have been formulated from blends of wheat/cashew kernel meals (Aroyeun, 2009), cassava flour/cashew apple powder (Ogunjobi and Ogunwolu, 2010), whole wheat and full fat soya (Ndife et al., 2014), wheat flour/cashew-apple residue as a source of fibre (Ebere et al., 2015), composite flour of plantain/Bambara groundnut protein concentrate (Kiin-Kabari and Giami, 2015).

Cashew (*Anacardium occidentale* Linn.) belonging to family *Anacardiaceae* is an extremely hardy tree that grows on poor soil under various climatic conditions. It is a native to Brazil and is being extensively grown in India, East Africa and Vietnam (Muniz et al., 2006). These countries including Nigeria are the main producers of cashew (Honorato et al., 2007). Cashew is one of the most important plantation crops earning huge amount of foreign exchange through its kernel and Cashew Nut Shell Liquid (CNSL). India is the World's largest exporter of cashew nuts followed by Brazil. Africa is third in terms of global production of cashew, producing approximately 100,000 tonnes of cashew nut per year. Africa gains little from the production as most of the nuts produced are mostly exported to the USA, Netherland and other European countries for processing. Some estimates suggested that African countries loose about one hundred million dollars per year by not processing their nuts which Nigeria is one of them (Ogunmoyela, 1983). Cashew consists mainly the nuts containing an embryo and a false fruit commonly called cashew kernel/nut and cashew-apple, respectively (Akinwale, 2000). In 2007, Nigeria produced 660,000 MT of raw nuts and given the weight ratio of apple to nut 8:1 (FAO, 2008; Cornier, 2008).

Cashew research started at the Cocoa Research Institute of Nigeria (CRIN), Ibadan in 1971 (Hammed et al., 2007). The study revealed that only 10% of cashew produced in Nigeria is utilized (Oduwole et al., 2001). The practical implication of this is that 90% is wasted annually. This led to the studies of the physicochemical characteristics of cashew nut flour (Aremu et al., 2006), cashew kernel meals enriched cookies (Aroyeun, 2009), roasted and defatted cashew nut flour (Omosuli et al., 2009), effect of packaging materials, storage time and temperature on the colour and sensory characteristics of cashew-apple juice (Emelike and Ebere, 2015) and composite flour of wheat and cashew-apple residue as a source of fibre in cookie preparation (Ebere et al., 2015). All these efforts were aimed at reducing the wastage of cashew in Nigeria and to use its by-products (cashew-apple residue and cashew nut flour) as functional ingredients.

The functional properties of commonly used plant materials like soybean, cowpea and pigeon pea are studied extensively by many scientists (Narayana and Rao 1982). To evaluate how their flour proteins can be used to supplement or replace more expensive protein sources which are used traditionally (Akobundu et al., 1982) and to upgrade a carbohydrate rich foods such as snack and pap. However, research on the functional potentials of the defatted and undefatted cashew kernel flour is few. Hence, the purpose of this research work is to evaluate the effect on the proximate, mineral and functional properties of defatted and undefatted cashew kernel flour.

MATERIALS AND METHODS

Collection of Materials

Mature, ripe cashew (*Anacardium occidentale* Linn.) apples, red and yellow varieties were harvested in an orchard at Uturu, Abia State, Nigeria. A total of ten kilograms of cashew nuts were used. All chemicals used were of analytical grade.

Preparation of Cashew Nut Flour

The nuts were sun dried for three days to prevent deterioration during storage. They were conditioned (mild spraying with water in a sieve) to increase flexibility and prevent scorching.

The conditioned cashew nuts were divided into portions of 500g each for easy processing, placed in a metal basket and immersed in a pot of hot vegetable oil (Corn oil) for 1 min to make the shell brittle for shelling. The nuts were stirred at intervals of 10 sec to prevent burning while in the hot oil. The Cashew Nut Shell Liquid (CNSL) of the nut extracted into the pot as the volume of the oil increases in the pot. The cashew nuts were poured out after 1 min and allowed to cool for about 1h. The brittle shell was broken with wooden mallet and the kernel extracted. The kernels were pooled together which yielded 6.57kg. The kernel was further roasted in hot sand for 30 min to facilitate easy removal of testa, cooled slightly and the testa removed manually using tooth pick.

Defatted and Undefatted Cashew Kernel Flour

The kernel was cleaned, sorted for rancid kernels, insect damage and discoloured kernels and divided into two equal parts. One was used to process undefatted cashew kernel flour while the other defatted cashew kernel flour. The kernels were milled separately using Kenwood blender (Model A907D U.K) for size reduction which formed oily cashew kernel flour. One part of the obtained flour was sieved using 0.400mm sieve and left at that stage as the undefatted cashew kernel flour. The other was further processed for the defatted flour by oven drying it at 105°C for 1h to reduce the moisture content and to agitate the fat molecules of the flour. The oil was extracted by solvent extraction with petroleum ether (b.p 40 – 60°C) in continuous soxhlet extraction apparatus for 3h and then wrapped in filter paper. The flour produced was sieved to remove larger particles or grains, forming a fine powder. The defatted and undefatted samples were used to analyse the proximate, mineral and functional properties.

DETERMINATION OF PROXIMATE COMPOSITIONS

The recommended methods of Association of Official Analytical Chemist (AOAC, 2012) were employed in determining the moisture, crude protein, ether extract, ash and crude fibre contents of defatted and undefatted cashew kernel flour. Carbohydrate was determined using the Clegg Anthrone reagent method as described by Osborne and Voogt (1978).

DETERMINATION OF MINERAL CONTENTS

Mineral elements of defatted and undefatted cashew kernel flour were analysed for; phosphorous, iron, calcium, sodium, potassium, magnesium and manganese using an Atomic Absorption Spectrophotometer, AAS (Model 372, Perkin – Elmer, Beaconsfield, U.K) by wet digestion according to the method described by AOAC (2012).

DETERMINATION OF FUNCTIONAL PROPERTIES

Water Absorption Capacity

Water absorption capacity of both samples was determined using Sosulski et al., (1979) method. One gram of cashew kernel flour was stirred into 10ml of distilled water in a centrifuge tube, after which the suspension was allowed to stand for 10 min and then centrifuged for 25 min at 2,500 rpm. The clear supernatant was decanted into a measuring cylinder and the adhering drops of water were removed by inverting the tube and the tube was weighed. The weight of water absorbed by 1.0g of flour was calculated and expressed as water absorption capacity.

$$\text{Water absorption capacity (\%)} = \frac{\text{final weight of } Y - X}{X} = 100$$

Where Y = Final weight of sample and tube and X = Initial weight of sample and tube.

Fat Absorption Capacity

The method of Lin et al., (1974) was used in the determination of fat absorption capacity. One gram of cashew kernel flour was added to 10ml of groundnut oil in a centrifuged tube, stirred together and allowed to stand for 30 min at room temperature (30±2°C). The sample was centrifuged at 3,000 rpm for 30 min. The clear supernatant was decanted into a measuring cylinder, the adhering oil was removed by inverting the tube and the tube was weighed. The weight of oil absorbed by 1.0g of flour was calculated and expressed as fat absorption capacity in percentage.

$$\% \text{ Fat absorption capacity} = \frac{Y - X}{X} \times \frac{100}{1}$$

Where Y = Final weight of sample and tube and X = Initial weight of sample and tube.

Bulk Density

Bulk density was determined on the defatted and undefatted cashew kernel flour using the method of Narayana and Rao (1984). A weighed centrifuge tube of 10ml was filled with 10g of cashew kernel flour and the weight was recorded. This was subjected to a constant tapping with subsequent addition of the flour until no further change in volume was observed and then final weight was taken. The difference in weight was used to calculate the bulk density of the samples.

$$\text{Bulk density} = \frac{\text{weight difference}}{\text{Initial volume}}$$

Foam Capacity and Stability

Foam capacity and stability of the samples were determined using Coffman and Garcia (1977) method with some modifications. Two grams of defatted and undefatted cashew kernel flour was whipped with 10ml distilled water for 5 min in a Kenwood blender (Model A907D U.K) at high speed and poured into 250ml graduated cylinder. The volume of foam at 30 sec after whipping was expressed as the foam capacity and the volume of the foam over 30 – 60 min as foam stability for the respective time periods. The volume increase (%) was calculated according to the following equation;

$$\begin{aligned} \text{Volume increase (\%)} \\ = \frac{\text{volume after whipping (ml)} - \text{volume before whipping (ml)}}{\text{Volume before whipping}} \times \frac{100}{1} \end{aligned}$$

Emulsification Capacity and Stability

Emulsion capacity and stability was studied using the method described by Beuchat (1977). Two grams (2g) of samples were blended in a Kenwood blender (Model A907D, U.K) with 100ml of distilled water for 30 sec at high speed. After complete dispersion, vegetable oil (Corn oil) was added continuously in 5ml portions from a burette. Blending continued until the

emulsion breakpoint (separation into two layers) was reached. The emulsion capacity was determined at room temperature ($30\pm 2^\circ\text{C}$) and the emulsion capacity was calculated using the formula;

$$\text{Emulsion capacity (\%)} = \frac{\text{Height of emulsion}}{\text{Total height of liquid}} \times \frac{100}{1}$$

The stability of the emulsions were evaluated by keeping the emulsion at room temperature ($30\pm 2^\circ\text{C}$) for 20h, noting the separation of water in the graduated cylinders, the emulsion stability was calculated using the equation;

$$\text{Emulsion stability (\%)} = \frac{\text{Volume of water separated}}{\text{Total volume of liquid}} \times \frac{100}{1}$$

Nitrogen Solubility Index

Nitrogen solubility of the flour samples were determined using the method described by Narayana and Rao (1984). Quantity of 0.5g of the defatted and undefatted cashew kernel flours was mixed with 30ml phosphate buffer between pH ranges of 8 – 9. The mixture was allowed to stand for 2h at room temperature ($30\pm 2^\circ\text{C}$). The suspension was then centrifuged for 40 min. At a speed of 2,000rpm, the supernatant was decanted and used to estimate the nitrogen content by Kjeldahl method. Nitrogen extracted was expressed as the percentage of the flour nitrogen.

$$\text{Nitrogen solubility index (\%)} = \frac{\text{Soluble nitrogen}}{\text{Total nitrogen of sample}} \times \frac{100}{1}$$

Statistical Analyses

Results were expressed as mean values and standard deviation of five (5) determinations. The obtained data were analysed using a one-way analyses of variance (ANOVA) using Statistical Packaging for Social Science (SPSS) version 20.0 software 2011 to test the level of significance ($p < 0.05$). Duncan New Multiple Range Test was used to separate the means where significant differences existed.

RESULTS AND DISCUSSIONS

Proximate Compositions

Moisture content of cashew kernel flour was 4.4% for defatted and 5.7 for undefatted as shown in Table 1. Hussain et al., (2008) reported a significant reduction in the moisture content of partially defatted flaxseed which agrees with the observation in this study. This indicates that defatting decreases the moisture content of raw materials and food materials with low moisture content have the benefit of been stored for a longer period. Protein content of the defatted flour (34.0%) is significantly high ($p < 0.05$) compared to that of undefatted sample (19.8%). This is within the range of protein content of soybean (35.1%) and melon seed (33.3%) reported by Achinewhu (1983) and Hussain et al., (2008) for partially defatted flaxseed flour (34.6%). It was significantly high to that reported by Omosuli et al., (2009) for roasted and defatted cashew nut flour (27.31%) and Aremu et al., (2006) for cashew nut flour (25.3%). The difference in these reports could be the type of processing methods applied which might have subdued the

protein content of the raw material. The recommended daily allowance of protein for children ranges from 23.0 – 36.0g and adults 44 – 56g (NRC, 1989). However, it can be evaluated that defatted cashew kernel flour is a good source of protein substitute/enrichment since it can supply the recommended daily intake of protein for children. Defatting significantly decreased the ether extract (crude fat) of cashew kernel flour with 1.6% for defatted and 47.1% for undefatted sample. These values are in agreement with those reported by Fetuga et al., (1974) for undefatted cashew kernel, melon seed and groundnut (48.1, 53.3 and 55.2, respectively) and are regarded as oil seeds (Achinewhu, 1983). Crude fibre of defatted flour (6.2%) is significantly different ($p<0.05$) compared to undefatted sample (1.2%) and it compared very well with those of legumes with mean values ranging from 5 to 6% as reported by Anonymous (1972) and Aremu et al., (2006). Okon (1983) reported that a diet low in fibre is undesirable as it could cause constipation and that such diets have been associated with diseases of the colon like piles, appendicitis and cancer. This is an indication that cashew kernel flour can be incorporated in the formulation of snack products. Ebere et al., (2015) reported an increase in fibre content of cookies formulated from the blends of wheat/cashew apple residue up to 9.4% with 20% incorporation. The consumption of a moderate amount of cashew kernel will suffice in providing a reasonable proportion of the daily dietary need of individuals. Ash content of the defatted sample (1.8%) was significantly lower ($p<0.05$) compared to the undefatted flour (4.7%). The value for ash content of undefatted cashew kernel flour falls within the range of those reported by Aremu et al., (2006) and Omusuli et al., (2009) for cashew nut flour (4.4%) and roasted/defatted cashew nut flour (4.41%), respectively. Pomeranz and Clifton (1981) have recommended that ash contents of nuts, seed and tubers should fall within the range of 1.5 – 2.2% in order to be suitable for animal feeds. Therefore, defatted cashew kernel flour with ash content of 1.8% can be used for this purpose. The carbohydrate value of the defatted flour (32.2%) is comparatively higher to that of undefatted flour (21.5%). This is an indication that both defatted and undefatted cashew kernel flour is a good source of energy and capable of supplying the daily energy requirement of the body. Defatting made all the proximate compositions of cashew kernel flour more assessable.

Table 1. Proximate compositions of defatted and undefatted cashew kernel flour.

Parameters (%)	Defatted flour	Undefatted flour
Moisture	4.4±0.15 ^b	5.7±0.24 ^a
Protein	34.0±0.01 ^a	19.8±0.02 ^b
Ether extract	1.6±0.02 ^b	47.1±0.01 ^a
Crude fibre	6.2±0.10 ^a	1.2±0.14 ^b
Total ash	1.8±0.02 ^b	4.7±0.10 ^a
Carbohydrate	32.2±0.10 ^a	21.5±0.03 ^b

Results represent mean ± standard deviation of five determinations.

Values not followed by the same superscript within the rows are significantly different ($p<0.05$).

Mineral Compositions

The result showed that the mineral content of the defatted cashew kernel flour are most rich in magnesium (220mg/100g) followed by calcium (75.3mg/100g), Potassium (52.0mg/100g) and phosphorous (24.7mg/100g) which was significantly higher ($p>0.05$) to undefatted cashew kernel flour as presented in Table 2. This result is in a reverse case for the one reported by Omusuli et al., (2009). They reported that the most abundant of the mineral compositions of

roasted and defatted cashew nut flour was potassium followed by magnesium then calcium and was in a close agreement with the observation of Aremu et al., (2006). Magnesium observed in this study as the richest mineral compared favourably with the values reported by Offem (1990) for melon seed, soybean and cocoa bean. Iron and sodium is the least abundant minerals in this study with the values of 11.8mg/100g and 7.8mg/100g, respectively. It was observed that all the analysed mineral components were significantly higher ($p < 0.05$) in defatted cashew kernel flour than the undefatted sample except manganese with significantly lower value in the defatted flour (2.9mg/100g). This observation is in agreement with the report of Hussain et al., (2008) who reported that partial defatting resulted in a significant increase in the mineral content of the flaxseed flours. This is an indication that fat content in cashew kernel and flaxseed flours is an anti-mineral element. Generally, defatted and undefatted cashew kernel flour is comparatively low in phosphorous, iron, sodium and manganese when compared to the values of melon seed and soybean but the values are high enough to contribute significantly to the Nigerian diet and thus mitigate the prevalent micronutrient deficiency disease.

Table 2. Mineral content of defatted and undefatted cashew kernel flour

Parameters (mg/100g)	Defatted flour	Undefatted flour
Phosphorous	24.7±0.20 ^a	8.3±0.03 ^b
Iron	11.8±0.15 ^a	9.0±0.12 ^b
Calcium	75.3±0.10 ^a	59.0±0.20 ^b
Sodium	7.8±0.03 ^a	5.0±0.01 ^b
Potassium	52.0±0.15 ^a	6.8±0.04 ^b
Magnesium	220±0.24 ^a	190±0.12 ^b
Manganese	2.9±0.07 ^b	9.0±0.14 ^a

Results represent mean ± standard deviation of five determinations.

Values not followed by the same superscript within the rows are significantly different ($p < 0.05$).

Functional Properties

The essentials in determining the potential uses for cashew kernel flour are its functional properties. The bulk density of the cashew kernel flour was highest with undefatted flour (0.2g/ml) and lowest with defatted flour (0.1g/ml) as shown in Table 3. From the result, it is obvious that defatting cashew kernel flour reduces its bulk density. This observation is in agreement with the result of Hussain et al., (2008); they reported that bulk density of the flours decreases as a result of defatting process. Bulk density plays an important role in packaging, transportation of food products and decreases porosity of materials due to surface properties (Milson & Kirk, 1980). Water absorption capacities for defatted flour (7.7%) was significantly higher than the undefatted flour (0.8%) and were very low when compared to the values of water absorption capacities of soybean (130%) and cowpea (100%) as reported by Lin et al., (1974). However, the defatted flour absorbed more water and fat than the undefatted flour. The fat absorption capacity for the defatted flour was significantly different ($p < 0.05$) than the undefatted sample with the values of 20.0% and 6.2%, respectively. The values reported in this work are lower than those reported for most nuts and seeds; pumpkin seed 87%, wheat flour 84.2% and soy flour 84.4% (Olaofe et al., 1994). It was high than the values of roasted full fat and roasted defatted (1.31 and 1.27g/g, respectively) for flaxseed flours as reported by Hussain et al., (2008) and gourd seed 6%. Defatted cashew kernel flour can be useful in production of low-fat bakery products and used as flavour carriers or modifiers in fabricated foods as meat

analogues. Good flavour carriers should bind flavours tightly, retain it during processing and release it during mastication of the food in the mouth.

The result showed that foam capacity (4.0ml) and stability (106ml) at 30 min was higher in defatted cashew kernel flour than the undefatted flour. This is similar to the report of Hussain et al., (2008) who reported decreased in roasted as compared to non-roasted flaxseed flours. Giami and Bekebian (1992) also reported that the foam of the defatted flour was more stable than that of the full fat samples. At 1h, the volume of foam stability both for defatted and undefatted (106 and 104 ml, respectively) remains constant as at 30 min. This is an indication that foam stability attained its maximum volume at 30 min. There was no significant difference ($p>0.05$) between the emulsion capacities of defatted and undefatted cashew kernel flour with the mean values of (5.0%). Emulsion stability at 30 min and 1h obtained the same value and defatted flour is reported to have high stability value than undefatted flour (88.0 and 84.0%), respectively. These values are higher than those reported for wheat and soybean flours (7.1 and 18.0%), respectively. This shows that cashew kernel flour can be substituted to soybean/wheat flours as meat extenders/binders in stabilizing colloidal fat and in turn reduce the importation of wheat flour into Nigerian country. The nitrogen solubility of cashew kernel flour was highest for defatted flour at pH₈ (0.6%) compared to undefatted flour (0.2%). At pH₉, both defatted and undefatted samples obtained constant nitrogen solubility value of 0.8%.

Table 3. Functional properties of defatted and undefatted cashew kernel flour

Parameters	Defatted flour	Undefatted flour
Bulk density (g/ml)	0.1±0.00 ^a	0.2±0.01 ^a
Water absorption capacity (%)	7.7±0.20 ^a	0.8±0.01 ^b
Fat absorption capacity (%)	20.0±0.50 ^a	6.2±0.03 ^b
Foam capacity (ml)	4.0±0.02 ^a	3.9±0.01 ^a
Foam stability (30 min)	106±0.70 ^a	104±0.60 ^a
Foam stability (1h)	106±0.50 ^a	104±0.50 ^a
Emulsion capacity (%)	5.0±0.04 ^a	5.0±0.04 ^a
Emulsion stability (30 min)	88.0±0.03 ^a	84.0±0.02 ^b
Emulsion stability (1h)	88.0±0.02 ^a	84.0±0.01 ^b
Nitrogen solubility (pH ₈)	0.6±0.01 ^a	0.2±0.01 ^b
Nitrogen solubility (pH ₉)	0.8±0.05 ^a	0.8±0.01 ^a

Results represent mean ± standard deviation of five determinations.

Values not followed by the same superscript within the rows are significantly different ($p<0.05$).

CONCLUSION

The importance in determining the potential uses for cashew kernel flour are its proximate, mineral and functional properties. From the result, it can be concluded that defatted cashew kernel flour is a good source of crude fibre and protein concentrate as it can meet up the recommended daily intake of protein for children. This indicates that the flour can serve as a functional ingredient in the formulation of snack products, be substituted to soybean since the fibre content compares very well with those of legumes/wheat flours and in turn reduce the importation of wheat flour into Nigeria. Defatted cashew kernel flour with ash content of 1.8%

falls within the recommended range for animal feed and can also be used for this purpose. Generally, it was observed that all the analysed proximate, mineral and functional properties were significantly high ($p < 0.05$) in defatted cashew kernel flour than the undefatted sample. Therefore, cashew kernel flour can be defatted and use as a source of raw material.

REFERENCES

- Achinewhu S.C (1983). The saponin content of some Nigerian oil seeds. *Quality Plant Foods and Human Nutrition*, 33: 3 – 9.
- Akinwale, T. D. (2000) Cashew apple juice. “It’s uses in fortifying the nutritional quality of some tropical fruits”. *European Food Research Technology*, 211; 205 – 207.
- Akobundu E.N.T, Cherry J.P and Simmons J.G (1982). Chemical, functional and nutritional properties of egusi seed protein products. *Journal of Food Science*, 47: 829 – 835.
- Anonymous (1972). Physicochemical characteristics, fatty acid composition and lipoxgenase activity of crude pumpkin and melon seed flour. *Journal of Agriculture and Food Chemistry*, 44: 964 – 966.
- AOAC (2012). Official methods of analysis of AOAC International (19th ed.), Gaithersburg, M.D. USA.
- Aremu M.O, Olonisakin A, Bako D.A and Madu P.O (2006). Compositional studies and physicochemical characteristics of cashew nut (*Anacardium occidentale*) flour. *Pakistan Journal of Nutrition*, 5(4), 328 – 333.
- Aroyeun S.O (2009). Utilization of cashew kernel meals in the nutritional enrichment of biscuit. *African Journal of Food Science*, 3(10), 316 – 319.
- Beuchat L.R (1977). Functional and electrophoretic activities of peanut flour. *Journal of Agriculture and Food Chemistry*, 25: 252.
- Coffman G.W and Garcia V.V (1977). Functional properties and amino acid content of protein isolate. *Journal of Food Technology*, 12: 473.
- Cornier R (2008). Clarification of cashew apple juice and commercial applications. Oxfarm Quebec, Benin, West Africa.
- Ebere C.O, Emelike N.J.T and Kiin-Kabari D.B (2015). Physico-chemical and sensory properties of cookies prepared from wheat flour and cashew-apple residue as a source of fibre. *Asia Journal of Agriculture and Food Science*, 3(2), 213-218.
- Emelike N.J.T and Ebere C.O (2015). Effect of packaging materials, storage time and temperature on the colour and sensory characteristics of cashew (*Anacardium occidentale* L.) apple juice. *Journal of Food and Nutrition Research*, 3(7), 410 – 414.
- FAO (2008). Agriculture, Food and Nutrition for Africa, Rome, 385-387.
- Fetuga B, Babatunde G and Oyenuga V (1974). Composition and nutritive value of cashew nut to the rat. *Journal of Agriculture and Food Chemistry*, 22: 678 – 682.
- Giami S.Y and Bekebian D.A (1992). Proximate composition and functional properties of raw and processed full fat fluted pumpkin (*Telfairia occidentalis*) seed flour. *Journal of Science of Food and Agriculture*, 59: 321 – 325.
- Hammed L.A, Adedeji A.R, Asogwa E.U and Ibremo O.S (2007). Constraints to cashew production in Nigeria. A paper presented at the cashew stake-holders meetings organised by the African Cashew Alliance (ACA) held at IITA, Ibadan Nigeria. January, 26, p.12.
- Honorato T.L, Rabelo M.C, Goncalve L.R.B, Pinto G.A.S and Rodrigue S (2007). Fermentation of cashew apple juice to produce high added value products. *World Journal of Microbial Biotechnology*, 23: 1409 – 1415.

- Hurs H. and Martins S. (2005). Low carbohydrate and beyond: The health benefits of Insulin. *Cereal Food World*, 50, 57 – 60.
- Hussain S, Anjun F.M, Butt M.S and Sheikh M.A (2008). Chemical compositions and functional properties of flaxseed flour. *Sarhad Journal of Agriculture*, 24(4), 649 – 653.
- Kiin-Kabari D.B & Giami S.Y (2015). Physico-chemical properties and in-vitro protein digestibility of non-wheat cookies prepared from Plantain flour and Bambara Groundnut Protein concentrate. *Journal of food research*, 4(20), 78 – 86.
- Lin R.U, Humbert E.S and Sosulki (1974). Certain functional properties of sunflower meal products. *Journal of Food Science*, 39: 368 – 370.
- Milson T.S and Kirk P.C (1980). Legumes in human nutrition. *Food and Agriculture Organisation Nutrition Studies*, 19: 223 – 235.
- Narayana K and Narasinga-Rao M.S (1982). Functional properties of raw and heat processed winged bean (*Psophocarpus tetragonolobus*) flour. *Journal of Food Science*, 42: 534 – 534.
- Narayana K and Narasinga-Rao M.S (1984). NISPRI, 1990. Fruits and Vegetable: Storing your Produce Advisory Booklet. *Journal of Food Science*, 47(4), 1534 – 1538.
- National Research Council (NRC, 1989). Recommended Dietary Allowances (10th edn). National Academic Press, Washington, DC, USA, 24.
- Ndife J, Kida F and Fagbemi S. (2014). Production and quality assessment of enriched cookies from whole wheat and full fat soya. *European Journal of Food Science and Technology*, 2(1), 19 – 28.
- Oduwole, O.O., Akinwale T.O. and Olubamiwa. O. (2001). Economic evaluation of a locally fabricated extraction machine for a cottage cashew juice factory. *Journal of Food Technology of Africa*. 6 (1); 18 – 20.
- Offem J.O (1990). Individual variation in amino acid and chemical composition of defatted cocoa bean meal from South Eastern Nigeria. *Journal of Science of Food and Agriculture*, 54: 129 – 135.
- Ogunjobi, M.A. & Ogunwolu, S.O. (2010). Physico-chemical and sensory properties of cassava flour biscuits supplemented with cashew apple powder. *Journal of Food Technology*, 8(1), 24-29.
- Ogunmoyela, A.O. (1983). Prospects of cashew apple processing and utilization in Nigeria. *Process Biochemistry*, 3: 6 - 7.
- Okon B.D (1983). Studies on the chemical composition and nutritive value of the fruits of African star apple. M.Sc. Thesis University of Calabar, 67.
- Olaofe O.F, Adeyemi O and Adediran G.O (1994). Amino acid and mineral composition and functional properties of some oil seeds. *Journal of agriculture and Food Chemistry*, 42: 878 – 884.
- Omosuli S.V, Ibrahim T.A and Oloye D. (2009). Proximate and mineral composition of roasted and defatted cashew nut flour. *Patistan Journal of Nutrition*, 8(10), 1649 – 1651.
- Osborne D.R and Voogt P (1978). The analysis of nutrients in foods. London Academic Press, 130 – 134.
- Pomeranz A and Clifton D (1981). Properties of defatted soybean, peanut, field and pecan flours. In: Food analysis theory and practices west port L.T. AVI Publishing Comp. 17. *Journal of Food Science* 42: 1440 – 1450.
- Rosengarten F (1984). The Book of Edible Nuts. 5th ed. Walker and Co New York, 45.
- Sosulski F.W, Garratt M.D and Slinkard A.E (1976). Functional properties of ten legume flours. *Institute of Food Science and Technology Journal*, 9: 66 – 69.
- Toyin A (2001). A review of the prospect of cashew industry, 39.

WHO/FAO. (2003). Diet, nutrition and the prevention of chronic diseases. Report of a WHO/FAO expert consultation, World Health Organisation Technical report series 916, WHO Geneva.