

QUALITATIVE PHYTOCHEMICAL ANALYSIS OF ETHANOL LEAVES EXTRACT OF *JATROPHA TANJORENSIS* AND ITS EFFECTS ON LIVER FUNCTION OF MALE ALBINO WISTAR RATS

Nse Udoka Ebe^{1*}, Grace. S. Effiong¹ and Aniebiet E. Ikot¹

¹Department of Biochemistry, Faculty of Basic Medical Sciences, University of Uyo, P.M.B. 1017, Uyo, Nigeria

ABSTRACT: *This research work focuses on the phytochemistry of *Jatropha tanjorensis* leaves extract, and its activities on specific serum liver enzymes. Phytochemical screening of the leaves extract was carried out following standard procedures. To check the effects of these phytochemicals on the serum liver enzymes biomarker (Aspartate amino transferase, Alanine amino transferase and Alkaline phosphatase), twenty (20) healthy and mature male albino wistar rats randomly distributed into four groups of five each were used. Group I was fed on water and normal rat feed, Groups II, III and IV were fed with water, normal rat feed and in addition, were administered *Jatropha tanjorensis* leaves extract 100mg/kg, 250mg/kg and 500mg/kg respectively for 14 days. The results of the phytochemical study revealed the presence of flavonoids, alkaloids, tannins, saponins, terpenoids, cardiac glycosides and anthroquinones in the extract. Similarly, treatment with the leaves extract showed a noticeable decrease in all the three serum liver enzymes investigated, AST: 355.40 ± 2.18 , 319.28 ± 2.18 , 255.14 ± 2.66 , ALT: 132.51 ± 1.29 , 138.44 ± 1.38 , 149.93 ± 1.95 , ALP: 116.32 ± 0.57 , 113.58 ± 1.77 , 104.81 ± 0.29 , as compared to the control group 410.24 ± 2.13 , 159.22 ± 0.85 and 120.84 ± 1.45 respectively. This finding suggests that *Jatropha tanjorensis* leaves exhibit hepato-protective properties which could aid in the prevention of liver damage.*

KEYWORDS: *Jatropha tanjorensis* leaves extract, Phytochemistry, Alanine amino transferase, Aspartate amino transferase, Alkaline phosphatase

INTRODUCTION

Plants have been used since the medieval times for medicinal purposes for the treatment of all kinds of ailments. Some plants apart from serving as food have also been known to exhibit medicinal properties. Over 400,000 species of tropical flowering plants have medicinal properties, (Akpulu *et al.*, 1994). An estimated 60% of antitumor and anti-infectious drugs already in the market or under clinical trials are of natural origins, (Sofowora, 2008).

The medicinal properties of these plants are suspected to be as a result of the antioxidant, antipyretic, antimicrobial effects of secondary metabolites of phytochemicals in them. (Soetan

and Aiyelaagbe, 2009). Although plant based natural medicines are popularly acclaimed to be safe, scientist advocate for proper toxicological studies (Oyewole *et al.*, 2008), in order to ensure safety in the use of natural medicines. Toxicity is the undesirable property of any drug or chemical capable of producing injurious or detrimental effects on a living organism. Whether or not these injuries occur depend on the amount of chemical absorbed. (Gossel and Bricker, 1990). The toxic effect caused by drugs is similar in man and other animals, hence the use of animal models in toxicological studies. Most toxic effects of drugs occur at a predictable time after administration. However, the target organ of toxicity is not necessarily the site of accumulation. (Oboh and Masodje, 2009). This study however does not include doses that can induce toxicity upon administration.

Jatropha tanjorensis belongs to the family Euphorbiaceae. It is commonly called Hospital Too Far, Catholic Vegetable (Iwalewa *et al.*, 2005). The leaf is commonly consumed in many parts of Southern Nigeria as it does not require any special conditions to grow in the Southern soil. It is popular as a medicinal plant and is commonly used as a natural remedy for diabetes, malaria, infection, hypertension and some other conditions in this region, (Olayiwola *et al.*, 2004). The roots of this plant have been documented to serve as booster of blood and immune system, (Ebe and Chukwuebuka, 2019).

From literature, much work has not been reported on the effect of ethanol leaves extract of *Jatropha tanjorensis* on liver function. This study therefore investigates phytochemical composition of *Jatropha tanjorensis* leaves qualitatively, and also assesses effect of ethanol extract of the leaves on liver function of male albino whistar rats. The liver functions as an essential organ making proteins and blood clotting factors, manufacturing triglycerols and cholesterol, glycogen synthesis and bile production. The effect of *Jatropha tanjorensis* leaves extract on the liver function of the experimental animals was determined by testing for liver enzymes such as Aspartate amino transferase (AST), Alanine amino transferase (ALT), and Alkaline phosphatase (ALP).

MATERIALS AND METHODS

Fresh leaves of *Jatropha tanjorensis* were collected from an uncultivated land in Uyo Local Government Area of Akwa Ibom State. The leaves were authenticated at the department of Botany University of Uyo, Uyo, Nigeria. The leaves were washed and air dried at room temperature in the laboratory for three (3) weeks and grinded into powder form using a mechanical grinder. 450g of this sample was soaked in 1.5 litres of 80% ethanol in a large container, stirred, covered and left for 24 hours. This extract was filtered and placed in a water bath for 72 hours at 50°C to get rid of the excess solvent (ethanol). The crude extract obtained was used for the research.

Qualitative Phytochemical Analysis of Extracts

Standard procedures were adopted for qualitative phytochemical analysis of the leaves extract using High performance Liquid Chromatography (HPLC) and spectrophotometry. In addition, the steps described by Sofowora (2008) were followed.

Test for saponins: In a clean autoclaved test tube, 10ml of sterile water was taken. Then 0.5 g of the ethanol extract was supplemented to the test tube containing distilled water. The test tube was stirred vigorously; foaming was formed in the test tube which was the indication of the presence of saponins in the ethanol extract of the *J. tanjorensis*.

Test for tannins and phenolic: A cleaned sterilized test tube was taken, 0.5 g of ethanol extract was supplemented to the test tube and 12 ml of sterile water was also supplemented to the test tube. It was stirred and then filtered. The filtrate was transferred to another test tube. Some droplets of 6% FeCl₃ was supplemented to the filtrate, blue black or blue green coloration or aggregation indicated the occurrence of phenolic and tannins in the ethanol extract of *J. tanjorensis*.

Test for steroids: A cleaned and dried test tube was taken, 0.5 g of the ethanol extract was supplemented along with 4 drops of acetic anhydride and a droplet of concentrated H₂SO₄ was also supplemented. The combination was allowed to stand for 1 hour and neutralized with NaOH, a blue green color indicated the presence of steroid in the ethanol extract.

Test for glycosides: A cleaned and dried test tube was taken; 0.5 g of ethanol extract was supplemented to the test tube along with 3 ml of chloroform. The extract softened in chloroform. Concentrated H₂SO₄ was carefully supplemented to the test tube. A reddish brown color at the border showed the existence of a steroidal molecule, which was a glycine portion of the cardiac glycosides.

Test for Flavonoids: In a cleaned and dried test tube 5 ml of ethanol extract was supplemented along with concentrated H₂SO₄ (1ml) and 0.5 g of Magnesium. A pink coloration that disappeared on standing for 3 minutes showed the presence of flavonoids.

Test for alkaloids: In a 100 ml cleaned flask, 20 ml ethanol solution was supplemented, the solvent was evaporated by heating the beaker. The dried residue obtained was dissolved in 5 ml of HCl (2N) and filtered. 3 drops of Meyers reagent was supplemented, the presence of precipitate indicated the alkaloids.

Test for phylobatannins: In a cleaned test tube 3 ml of ethanol extract was supplemented along with 3 ml of 2% HCl and the combination was allowed to bioleach. Appearance of a red aggregation was taken as an evidence for the occurrence of phylobatannins in the ethanol extract.

Test for anthraquinones: 3 ml of ethanol extract was shaken and filtered and 6 ml of 12% ammonia was supplemented to the filtrate. The combination was allowed to stand and the presence of a pink color in the ammonical segment showed the existence of anthraquinones.

Test for terpenoids: In a cleaned test tube 3 ml of the ethanol extract was dissolved in 3 ml of chloroform and vaporized to dryness. 3 ml of concentrated H₂SO₄ was then supplemented and heated for 3 minutes, grayish color showed the existence of terpenoids

Experimental Animals and Extract Administration

Twenty (20) healthy and matured male albino wistar rats weighing between 150-262g were used in this study. The rats were housed in the Basic Medical Sciences animal house University of Uyo in wooden cages covered with net in a well-ventilated room under standard laboratory conditions. During the experimental period, animal feed (growers) and water were provided *ad libitum*. The animals were assigned into four (4) groups of five rats each. Group I served as the control group, Groups II, III and IV were administered *Jatropha tanjorensis* leaves extract 100mg/kg body weight, 250mg/kg body weight and 500mg/kg body respectively for 14 days. Administration was done orally using a fabricated cannula (constructed from syringe and feeding tube) and the doses were calculated based on the body weight of the animals. All experimental animals were treated in line with the guidelines of Institutional Animal Ethical Committee. At the end of the administration, the rats were fasted overnight (12 hours) and anaesthetized by putting each in glass chamber (desiccator) containing chloroform. The blood samples were obtained by cardiac puncture using sterile syringes and needles and placed into sterile plain tubes.

Sera samples were obtained from the clotted blood after centrifugation at 2000 rpm for 5 minutes and the supernatant serum was collected and stored in sample tubes and refrigerated for subsequent analysis.

Enzymes Assayed

Three enzymes assayed in this study were Aspartate amino transferase or Glutamic oxaloacetic transaminase (AST or GOT), Alanine amino transferase or Glutamic pyruvic transaminase (ALT or GTP), and Alkaline phosphatase (ALP) based on UV- kinetic method. All chemical kits used were of analytical grade, obtained from Randox Laboratories Limited, 55 Diamond road, Crumlin, Country Antrim, BT29 4QY, United Kingdom.

Estimation of Aspartate Amino Transferase (AST) Activity


Randox laboratories limited concentration reagents used were;

- (i) phosphate buffer (100mmol/l, pH 7.4,
- (ii) L-aspartate (100mmol/l),
- (iii) α -oxoglutarate (2 mmol/l),

(iv) 2,4-dinitrophenylhydrazine (2mmol/l).

Principle

The enzymatic reaction sequence employed in the assay of aspartate amino transferase is as follows:



Oxaloacetate formed in the first reaction is reduced to malate in the presence of malate dehydrogenase and NADH as cofactor. The activity of aspartate amino transferase is determined by measuring the rate of oxidation of NADH at 546nm.

Procedure

Twenty test tubes were secured in the test tube rack and 100µl of the test samples was pipetted into the test tubes. Five hundred (500) µl of buffer (reagent 1) was pipetted into the reagent blank and the test tubes. One hundred (100) µl of distilled water was pipetted into reagent blank. The mixture was mixed gently for 10 seconds and incubated for 30 minutes at 37°C. Thereafter, 500µl of 2,4-dinitrophenyl hydrazine was pipetted into the reagent blank and test tube samples. The mixture was mixed gently and allowed to stand for 20minutes at 25°C. Five thousand (5000) µl of Sodium Hydroxide was pipetted into the reagent blank and test tube samples. The mixture was mixed gently for 10 seconds. The absorbance of the sample was read against the reagent blank. The wavelength was set at 546nm and the reagent blank was used to zero the spectrophotometer. Each test sample was transferred carefully into cuvette and the absorbance was read against the reagent blank.


Estimation of Alanine Amino Transferase (ALT) Activity

Randox laboratories limited concentration reagent contents used were;

- (i) Phosphate buffer (100 mmol/l. pH 7.4),
- (ii) L- Alanine (200 mol/l).

Principle

The enzymatic reaction sequence employed in the assay of ALT is as follows:



Alanine amino transferase (ALT) catalyses the transfer of amino group from alanine to 2-oxoglutarate, forming pyruvate and glutamate. The pyruvate formed in the first reaction is reduced to lactate in the presence of lactate dehydrogenase and NADH (cofactor). The activity of alanine amino transferase is determined by measuring the rate of oxidation of NADH at 564 nm. ALT is measured by monitoring the concentration of pyruvate hydrazine formed with 2, 4-dinitrophenyl hydrazine.

Procedure

Twenty test tubes were secured in the test tube rack and 100µl of the test samples was pipetted into test tubes. Five hundred (500) µl of buffer (reagent 1) was pipetted into the reagent blank and the test tubes. One hundred (100) µl of distilled water was pipetted into reagent blank. The mixture was mixed gently for 10 seconds and incubated for 30 minutes at 37°C. Thereafter, 500µl of 2,4-dinitrophenyl hydrazine was pipetted into the reagent blank and test tube sample. This was mixed and allowed to stand for exactly 20 minutes at 25°C. Five thousand (5000) µl of sodium hydroxide was pipettes into the reagent blank and test tube samples. The mixture was mixed gently for 10 seconds. The wavelength was set at 546nm and the reagent blank was used to zero the spectrophotometer. Each test sample was transferred carefully into cuvette and the absorbance was read against the reagent blank.

Estimation of Alkaline Phosphatase Activity (ALP)


Alkaline phosphatase activity in serum was quantitatively determined by the use of Randox Laboratory reagent kit which employs the Kinetic procedure for the determination of enzymes activity.

Randox laboratory Limited concentration reagents used were:

- (i) Diethaniline buffer (1mol/l, pH 9.5),
- (ii) MgCl₂ (0.5 mmol/l),
- (iii) p-nitrophenylphosphate (10 mmol/l)

Principle

The enzymes sequence employed in the assay of alkaline phosphatase involves the reaction steps below.



The enzymes rate is determined using a 2-amino-2-methyl-1-propanol (AMP) buffer to measure activity of alkaline phosphatase in serum. Here, alkaline phosphatase catalyses the hydrolysis of colourless organic phosphate ester substrate, P-nitrophenyl phosphate, to yellow coloured product, P-nitrophenol and phosphate. The system monitors the rate of change of absorbance at 405 nm over a fixed time interval. This rate of change in absorbance is directly proportional to the alkaline phosphatase activity in serum.

Procedure

Twenty test tubes were secured in the test tube rack and 100 µl of the test samples was pipetted into the test tubes. Two hundred and fifty (250) µl of buffer (reagent 1) was pipetted into the reagent blank and the test tubes. One hundred (100) µl of distilled water was pipetted into reagent blank. The mixture was mixed gently for 10 seconds to ensure a complete mixture. Thereafter, 250 µl of P-nitrophenyl phosphate was pipetted into the reagent blank and test tube samples. The mixture was maintained at 37°C for 60 seconds. The mixture was mixed gently for 10 seconds. The wavelength was set at 405 nm using an empty cuvette with 1 cm light path to zero the spectrophotometer. Each test sample was transferred carefully into cuvette and the initial absorbance was read. The timer was started simultaneously and absorbance was read again after 1, 2 and 3 minutes. The average absorbance was calculated and used to calculate the concentration. This was done by multiplying the average absorbance by 2760 to calculate U/I of alkaline phosphatase activity.

Statistical Analysis

Data were analyzed using mean ± standard deviation, and comparison between the test groups and control group was performed using student's T' test. Means were considered significantly different at $p < 0.05$.

RESULTS

Table 1 below shows the results of the phytochemical screening of the ethanol extract of *Jatropha tanjorensis* leaves. Phytochemical analysis of the extract revealed the presence of flavonoids, alkaloids, tannins, saponins, terpenoids, cardiac glycosides and anthroquinones. Phylobatannins and steroids were absent.

TABLE 1: PHYTOCHEMICAL SCREENING OF ETHANOL LEAVES EXTRACT OF *Jatropha tanjorensis*

PHYTOCHEMICALS	INDICATIONS
Flavonoids	+
Alkaloids	+
Tannins	+
Phylobatannins	-
Steroids	-
Terpenoids	+
Saponins	+
Cardiac glycosides	+
Anthroquinones	+

Key:

+ = present

- = absent

The biochemical assessment of the liver enzymes of albino wistar rats treated with ethanol extract of *Jatropha tanjorensis* leaves are presented on table 2. The results obtained for AST were 410.24 ± 2.13 , 355.40 ± 2.18 , 319.28 ± 2.18 and 255.14 ± 2.66 for groups I (control), II, III and IV respectively. For ALT determination, the results showed 159.22 ± 0.85 , 132.51 ± 1.29 , 138.44 ± 1.38 and 149.93 ± 1.95 for groups I, II, III and IV respectively. For ALP determination, the results gave 120.84 ± 1.45 , 116.32 ± 0.57 , 113.58 ± 1.77 and 104.81 ± 0.29 for groups I, II, III and IV respectively.

TABLE 2: EFFECT OF ETHANOL LEAVES EXTRACT OF *Jatropha tanjorensis* ON LIVER FUNCTION OF MALE ALBINO WISTAR RATS

	Groups	Concentration mg/kg body weight	AST (μ l)	ALT (μ l)	ALP (μ l)
I	Control	410.24 ± 2.13	159.22 ± 0.85	120.84 ± 1.45	
II	100	355.40 ± 2.18	132.51 ± 1.29	116.32 ± 0.57	
III	250	319.28 ± 2.18	138.44 ± 1.38	113.58 ± 1.77	
IV	500	255.14 ± 2.66	149.93 ± 1.95	104.81 ± 0.29	

Values are given as Mean \pm Standard Deviation (SD), n=5.

Values are significantly different from Control at $p < 0.05$.

DISCUSSION

This study was conducted to determine the phytochemical composition of ethanol extract of *Jatropha tanjorensis* leaves and its effect on liver function of male albino wistar rats. The results indicated the presence of flavonoids, alkaloids, tannins, saponins, terpenoids, cardiac glycosides and anthroquinones in the extract. This finding is consistent with previous work by Ehimwenma and Osagie (2007).

Flavonoids have antioxidant activity which has many beneficial effects on the cardiovascular system, also, epidemiological studies have illustrated that heart diseases are inversely related to flavonoid intake ((Evans, 1989).). This implies that taking flavonoid containing foods reduced the chances of developing heart disease. Flavonoids prevent the oxidation of low-density lipoprotein, lowers the blood levels of cholesterol and triglycerides thereby reducing the risk for the development of atherosclerosis (Subramani and Casimir, 2002). Flavonoids have also been reported to have vaso-dilatory and inhibitory effects on platelet aggregation thereby preventing coronary heart diseases (Okwu, 2004).

Saponins have been reported to have beneficial effects on blood cholesterol levels by preventing its re-absorption. The non-sugar part of saponins also has antioxidant activity which may help to reduce risk of heart diseases (Oakenfull and Sidhu, 1990). Cardiac glycosides which were also found in the ethanol extract of *Jatropha tanjorensis* leaves have been variously used from time immemorial as diuretics and heart tonics due to their beneficial effects on the heart. Cardiac steroids are widely used in the modern treatment of congestive heart failure and for treatment of atrial fibrillation. They increase the force of contraction of the heart and are very useful for heart failure patients. They act by affecting the availability of intracellular Ca^{2+} for myocardial contraction or increasing the sensitivity of myocardial contractile proteins (Walker *et al.*, 2002).

Tannins have been reported to be responsible for decreases in feed intake, growth rate, feed efficiency, net metabolisable energy, and protein digestibility in experimental animals. Therefore, foods rich in tannins are considered to be of low nutritional value. However, it has been suggested that consumption of tannin containing beverages, especially green teas and red wines can cure or prevent a variety of illness including heart related diseases (Van-Burden and Robinson, 1981). Terpenoids occur in nearly every natural food (including *Jatropha tanjorensis*) and have been associated with protection from and treatment of heart disease due to their antioxidant properties (Wagner and Elmadfa, 2003).

Liver function test (LFT) is commonly used in clinical practices to screen for liver diseases, monitor the progress of a known disease and determine the effects of potentially hepatotoxic drugs (Dufour *et al.*, 2000). It also helps in detecting hepatocellular necrosis but is considered a less specific biomarker enzyme for hepatocellular injury (Ozer *et al.*, 2008) as it can also signify abnormalities in heart, muscle, brain or kidney (Dufour *et al.*, 2000). Data accumulated from this study showed that the ethanol leaves extract of *Jatropha tanjorensis* showed a significant

decrease ($P < 0.05$) in serum AST in groups II, III and IV as compared to the control. The ratio of serum AST to ALT can be used to differentiate liver damage from other organ damage (Nathwani *et al.*, 2005). Inflamed or injured liver cells release higher than normal amounts of specific chemicals including liver enzymes into the blood stream resulting in elevated levels of these liver enzymes after serum examination. The extracts of *Jatropha tanjorensis* leaves showed a significant decrease on the serum levels of AST, indicating that the leaves extract has no negative or toxic interaction on the hepatocytes. ALT activity in the liver is much higher than that of the serum. This implies that in the case of hepatocellular injury or death, release of ALT from the damaged liver cells increases measure ALT in the serum. Elevated level of this enzyme is released during liver damage. The estimation of this enzyme is a more specific test for detecting liver abnormalities since it is primarily found in the liver (Dufour *et al.*, 2000). However, lower enzymatic activities are also found in skeletal muscles and heart tissue. This enzyme detects hepatocellular necrosis. An elevated ALT level is therefore an indicator of hepatocellular damage and has been used as a marker of liver injury. From the results, it was discovered that ALT decreased significantly ($P < 0.05$) in groups II, III and IV when compared to the control. ALP is another biomarker employed to assess liver function. It is often employed to assess the integrity of the plasma membrane. ALP also may be elevated if bile excretion is inhibited by liver damage. Hepatotoxicity leads to elevation of the normal values due to the body's inability to excrete it through bile due to the congestion or obstruction of the biliary tract. In humans, increased ALP levels have been associated with drug induced cholestasis (Dufour *et al.*, 2000).

The decrease in liver enzymes concentration in the serum of the experimental rats studied is an indication that administration of *Jatropha tanjorensis* leaves extract may not cause liver damage. This result shows no apparent toxic effects of the extract on the liver of the experimental animals and could therefore be effective as a hepatoprotective agent.

CONCLUSION

Phytochemical screening of *Jatropha tanjorensis* leaves revealed the presence of bioactive compounds such as flavonoids, alkaloids, tannins, saponins, terpenoids, cardiac glycosides and anthroquinones.

This research also showed that the plant leaves extract was able to alter the concentration of certain liver enzymes in the serum, leading to the decrease of AST, ALT and ALP in the serum of the albino wistar rats. The results proved that the plant leaves extract could reduce the risk of liver damage or disease.

REFERENCES

- Akpulu, I. N., Dada, J. D., Odama, E. and Galadima, A. (1994). Antimicrobial Activity of Aqueous Extracts of some Nigerian Medicinal Plants. *Nigeria Journal of Botany*, 7(4): 45–48.

- Dufour, D. R., Lott J. A., Nolte, F. S., Gretch, D. R., and Koff, R. S. (2000) Diagnosis and monitoring of hepatic injury II: Recommendations for use of laboratory tests in screening, diagnosis and monitoring. *Clinical Chemistry*. **46**(1): 2050-2068.
- Ebe, N. U. and Chukwuebuka, E. V. (2019). Effect of ethanol roots extract of *Jatropha tanjorensis* on haematological indices of male albino wistar rats. *European Journal of Biology and Medical Science Research*, **7** (1):7-14.
- Ehimwenma, S.O. and Osagie, A. U. (2007). Phytochemical Screening and Anti-anaemic Effects of *Jatropha tanjorensis* Leaf in Protein Malnourished Rats. *Plant Archive*. **7**(6): 509-516.
- Evans, W.C. (1989). Pharmacognosy 2nd Edition. London. Balliere Tindall. pp. 167-235.
- Gossel, T.A. and Bricker, J.D. (1990). Principles of Clinical Toxicology. 2nd Edition; New York, Raven Press Ltd, USA. pp. 413.
- Iwalewa E.O., Adewumi C.O., Omisore N.O., Adebajji A.O., Azike C.K., Adigun A.O., Adesina O.A., Olowoyo O.G. (2005). Pro- and antioxidant effects and cytoprotective potentials of nine edible vegetables in southwest Nigeria. *Journal of Medicinal Food* **8** (4):539-544.
- Nathwani, R. A., Pais, S., Reynolds, T. B., and Kaplowitz, N. (2005) Serum alanine aminotransferase in skeletal muscle diseases. *Journal of Hepatology* **41**(2): 380-382.
- Oakenfull, D., and Sidhu, G. (1990). Saponins - A useful treatment for Hypercholesterolaemia. *European Journal of Clinical Nutrition*, **44**(5): 79-88.
- Oboh, F.O. and Masodje, H.I. (2009). Nutritional and Antimicrobial Properties of *Jatropha tanjorensis* Leaves. *American-European Journal of Scientific Research* **4**(1):7-10.
- Okwu, D. E. (2004). Phytochemicals and vitamin content of indigenous spices of South Eastern Nigeria. *Journal of Sustainable Agriculture Environment*. **6**(4): 30-34.
- Olayiwola, G., Iwalewa, E. O., Omobuwajo, O. R., Adeniyi, A. A., and Verspohi, E.J. (2004). The Antidiabetic Potential of *Jatropha tanjorensis*. *African Journal of Pharmacology* **89**(5):42-48.
- Oyewole I. O., Ibidapo, C. A., Moronkola, D. O., Oduola, A. O., Adeoye G. O., Anyasor G. N. and Obansa J. A., (2008). Anti-malarial and Repellent Activities of *Tithonia diversifolia* (Hemlock) Leaf Extracts. *Health Journal* **8**(6): 13-16.
- Oyewole, I.O., Magaji, Z.J. and Awoyinka, O.A. (2007). Biochemical and Toxicological Studies of Aqueous Extract of *Tithonia diversifolia* Leaves in Wistar Albino Rats. *Journal of Medicinal Plants Research* **1**(2):030-033.
- Ozer, J., Ratner, M., Shaw, M., Bailey, W., and Schomaker, S. (2008). The Current State of Serum Biomarkers of Hepatotoxicity. *Journal of Toxicology* **245**(7): 194-205.
- Soetan, K. O. and Aiyelaagbe, O. O. (2009). The Need for Bioactivity-Safety Evaluation and Conservation of Medicinal Plants - A Review by Department of Veterinary Physiology, Chemistry, Biochemistry and Pharmacology, University of Ibadan, Nigeria. *Journal of Medicinal Plants* **15** (7):314-324.
- Sofowora, A. (2008). Medicinal plants and traditional medicine in Africa. Ibadan. Spectrum Books Limited. pp 250-260.

- Subramani, S., and Casimir, C. A. (2002). Flavonoids and Antioxidant Activity of Georgia Grown *Vidalia* Onions. *Journal of Agricultural and Food Chemistry*, **50**(19): 5338-5342.
- Van-Burden, T. P. and Robinson, W.C. (1981). Formation of complexes between protein and Tannin. *Journal of Agriculture and Food Chemistry* **1**(5): 77-82.
- Wagner, K. H., and Elmadfa, I. (2003). Biological Relevance of Terpenoids. Overview Focusing on Mono-, Di- And Tetra-terpenes. *Annual Journal of Nutrition and Metabolism*. **47**(3): 95-106.
- Walker, A. F., Marakis, G., Morris, A. P. and Robinson, P. A. (2002). Promising Hypotensive Effect of Hawthorn Extract: A Randomized Double-Blind Pilot Study of Mild, Essential Hypertension. *Phytotherapeutics Research* **16**(1): 48-54.