#### SELECTIVE TRANSFORMATION OF MENTHONE FROM THE ESSENTIAL OIL OF MENTHA PIPERATA L. INTO MENTHONE 2,4 DINITROPHENYLHYDRAZONE AND UV-VISIBLE SPECTROPHOTOMETRIC ANALYSIS

# \*NDZELI LIKIBI Belline<sup>1</sup>, LOUBAKI Laurent<sup>2</sup>, NSIKABAKA Samuel<sup>1</sup>, OUAMBA Jean-Maurille<sup>1</sup>

<sup>1</sup>Plant and Life Chemistry Unit, Faculty of Sciences and Techniques, Marien Ngouabi University, Brazzaville, BP 69, Congo.

<sup>2</sup>Multidisciplinary Research Team in Food and Nutrition, Faculty of Science and Technology, University Marien Ngouabi, Brazzaville, BP 69, Congo.

**ABSTRAT:** The essential oil of Mentha piperata L, brown in color, was extracted by hydrodistillation from leaf stems collected in the southern district of Brazzaville (Bacongo), with a yield of 0.42 %. Analysis by gas chromatography (GC) and by gas chromatography coupled with mass spectrometry (GC/MS) revealed the presence of thirty (30) constituents of which twenty (20) were identified representing a proportion of (93.26 %) of the chemical composition of the total essential oil. The essential oil consists mainly of monoterpenes (92.47 %) with a predominance of oxygenated monoterpenes (89.43 %) dominated by menthol (44.08 %), followed by menthone (14.49 %), menthyl acetate (13.54 %) ), mint furanone 1 (7.81 %) and menthofuran (3.04 %). Hydrocarbon monoterpenes occupy a low level of (3.04 %) characterized by limonene (2.71 %), orthocymene (0.37 %),  $\beta$ -pinene (0.35 %) and  $\alpha$ -pinene (0.15 %). Concerning the group of sesquiterpenes, this represents a minimal rate (0.79 %), oxygenated compound are predominant (0.57 %) marked by caryophyllene oxide (0.40 %) and viridiflorol (0.17 %). We note the presence of a single hydrocarbon sesquiterpene, E caryophyllene occupying a small proportion of (0.22 %). Menthone 2,4-dinitrophenylhydrazone was hemi-synthesized by reacting the essential oil of Mentha piperata with 2,4dinitrophenylhydrazine in the presence of sulfuric acid, a simple, easy method occurring in a rapid time (five minutes ). The rate of conversion of menthone to menthone 2,4dinitrophenylhydrazone of orange color is 29 %. Analysis of menthone 2,4dinitrophenylhydrazone by UV-visible spectrophotometry showed an absorption maximum wavelength of 356 nm. The UV-visible spectrophotometric method employed for the determination of this hydrazone is convenient, fast and simple. The hemi-synthesized menthone 2,4-dinitrophenylhydrazone could be useful in the pharmaceutical, agrochemical, perfume and *cosmetic industries* 

**KEYWORDS:** Mentha piperata L., essential oil, menthone, menthone 2,4dinitrophenylhydrazone, UV-visible spectrophotometry, wavelength.

# **INTRODUCTION**

Essential oils contain a diverse range of organic compounds, including ketones. Among these is menthone; one of the main constituents of essential oils of certain species of the genus Mentha belonging to the family Labieaes or Lamiaceaes [1] including *Mentha piperata* L. with considerable proportions [2; 3; 4].

It is a cyclic ketone (carbonyl compound), more precisely a cyclohexanone substituted by isopropyl and methyl groups in positions 2 and 5 respectively. It is of great interest in cosmetics, perfumery, agrifood and pharmacy. Indeed, it enters into the composition of natural perfumes and aromas [5; 6]. It has various chemical properties due to the presence of the C=O double bond giving rise to several nucleophilic and electrophilic addition reactions [7]. However the main or disadvantage of this is its instability which often results in oxidations, hence the need to convert it into hydrazones, crystalline compounds more stable than their precursors (carbonyls) [8; 9].

Hydrazones are organic compounds of the general formula: RR'C = N-N-R''R ''.

2,4-Dinitrophenylhydrazones are substituted hydrazones, derived from the condensation of an aldehyde or a ketone with 2,4-dinitrophenylhydrazine [10] (figure 1); a reversible reaction subjected to acid catalysis [11]. They play an important role in the protection of carbonyl compounds [12]. They are also used for the isolation, purification and characterization of the carbonyl group [13] and as an intermediate in organic synthesis [14]. Furthermore, hydrazones are used for the extraction or determination of transition metals such as iron [15], molybdenum [16], by formation of hydrazonic metal complexes.

Menthone 2,4-dinitrophenylhydrazone, for its part, results from the condensation of menthone (a cyclic monoterpene ketone) with 2,4-dinitrophenylhydrazine.

In pharmacology, numerous studies carried out affirm that menthone hydrazone exhibits various biological activities, among others, anticonvulsant [17;18] antibacterial [19]; antimicrobial and cytotoxic [20]; antifungal [21]; anticancer; antidepressive; analgesic and antiviral [22].

In addition, it should be noted that hydrazide hydrazone derived from cyclohexanan-1,3-dione is endowed with anticancer activity [23].

Furthermore, thiosemicarbazone and semicarbazone derived from menthone have shown significant anti-HIV and analgesic activities [24, 25].

Also, menthopyrazole derived from menthone is endowed with electrochemical; inter alia, anticorrosive properties [26]. While hydrazone derived from formylmenthone has electrochemical complexing properties towards metals such as Cu (II) and Ni (II) [27].

Several synthesis of 2,4-dinitrophenylhydrazones from menthone have been reported in the literature [28; 29; 21; 30; 31; 32].

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

Very few studies of the hemi-synthesis of hydrazones derived from essential oil ketones are described in the literature.

To our knowledge, the hemi-synthesis of menthone 2,4-dinitrophenylhydrazone from essential oils are not known.

Various analytical methods are used for the determination of hydrazone ketones, among them, HPLC-UV is the technique of choice for the determination of the latter [33]. To this are added HPLC-MS [34], RRLC-UV and RRLC-MS (/ MS) | 35], LC-MS [36], as well as GC-MS [37; 38]. In addition, APPI-MS [39] and HPLC [40; 41] are also involved in the analysis of 2, 4-dinitrophenylhydrazones ketones.

Besides these methods, UV-visible spectrophotometry also is widely used for the determination of hydrazone ketones [42; 43; 44]. However, some of these analytical methods involve long analysis time, tedious sample pre-treatment, and high product cost.

The aim of this present work is to extract the essential oil of *Mentha piperata*, to make the hemisynthesis and to characterize the menthone 2,4-dinitrophenylhydrazone by UV-Visible spectrophotometry, a simple and fast method.



Figure 1: Reaction of a carbonyl compound with 2,4-dinitrophénylhydrazine.

# MATERIALS AND METHODS

#### Vegetable material

The samples of *Mentha piperata* (figure 2) were collected in april 2019 in the southern districts of Brazzaville (Bacongo). They were identified by botanists from the National Herbarium of the Congo. Only leaf stems were selected for the study.



Figure 2: Leaves of Menta piperata L.

## **Extraction of essential oils:**

After eight (8) days of drying at room temperature, in a ventilated room, the samples of *Mentha piperata* made up of the leaf stems are subjected to hydrodistillation for four (4) hours using a Clevenger type extractor [45], fitted with a two (2) liters balloon. The condensate loaded with essential oil and hydrolate is collected. The essential oil is separated from the hydrolate by decantation. The extraction with diethyl ether is carried out to isolate the aqueous phase of the essential oil followed by drying the ether phase with anhydrous sodium sulfate. Twenty-four (24) hours after evaporation of the diethyl ether in air, the essential oil is recovered. These operating conditions are summarized in Table I. The yield Y in essential oil is calculated according to the following formula:

 $Y = \frac{\text{Masse d'huile essential (g)}}{\text{Masse of plant material used (g)}} \ge 100$ 

Plant material	Mentha piperata
Organs	Leaf stems
Quantity of matter	131
dry (g)	
Quantity of water (L)	1
Execution time	4
of operation (h)	

Table I: Operating conditions for hydrodistillation of Mentha piperata L.

# **Organoleptic characteristics**

The organoleptic properties (appearance, colour, smell) were evaluated based on the sensory method involving sight and smell.

# Physicochemical parameters Refractive index:

The refractive index is the ratio between the sine of the angles of incidence and refraction of a light ray of a given wavelength, passing from the air in the essential oil maintained at a constant temperature [46]. It is linked to the chemical function to which the essential oil belongs and also depends on the acidity, the polymerization of the essential oil.The measurement was carried out using a Novex Abbé refractometer. The method described by AFNOR was applied [46]. It consisted of putting two (2) to three (3) drops of essential oil on the prism of the refractometer. Look through the eyepiece by rotating the adjustment knob until the dividing line of the bright and dark areas is in the center of the reticle. Read the refractive index value of the essential oil on the reading scale. The index is given by reading on the refractometer at the temperature T at which the reading is taken. This is brought to 20 °C according to the following formula:

n20 = nT + 0.00045 (T-20)

With T: temperature on reading is taken

#### Acid index:

The acid Index (Ia) is the number of milligrams of potash (KOH) required to neutralize the free acids contained in 1 gram of essential oil [46]. It makes it possible to check the quality of an essential oil, in particular its deterioration and aging during storage time. The method described by AOAC [47] was applied to determine the refractive index. It consists of placing 1 gram of essential oil in an Erlenmeyer flask in which 5 milliliters (mL) of 95 % ethanol and 5 drops of phenolphthalein are added. The mixture is warmed in a water bath to about 65 °C for 10 minutes. After cooling, it is titrated with a 0.1 N solution of potassium hydroxide (KOH) using a burette, until the solution turns pink. The operating conditions are summarized in Table II. Thus, the acid index is determined by the following formula:

Ia = 0.56 x V/m in which: V: volume of the KOH solution. m: mass of essential oil in gram

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

**Table II:** Operating conditions for determining the acid index of essential oil of *Mentha* piperata L.

Parameter	Values
Mass of essential oil in (g)	1
KOH cencentration (mol/L)	0.1
Volume of ethanol (mL)	5
Amount of phenolphthalein (drops)	5

#### Analysis of essential oils

#### Analysis by gas chromatography

Quantification of the constituents was performed using a Hewlett Packard HP 5890 type chromatograph equipped with a flame ionization detector fitted with HP ChemStation data acquisition software. The different constituents are separated using a DB5 capillary column (30 mx 0.25 mm), (film thickness 0.25  $\mu$ m) under the following operating conditions: helium carrier gas (1 mL.min-1), temperature injector temperature: 280 °C, detector temperature: 280 °C. The oven is programmed at 50 ° C for 5 minutes with a gradient of 5 °C.min-1 from 50 to 300 °C, 5 minutes at 300 °C with a split mode injection of 1-20.

#### Analysis by gas chromatography-mass spectrometry

Analysis by gas chromatography-mass spectrometry was carried out using a Hewlett Packard HP 6890 brand chromatograph coupled to an HP 5973 mass spectrometer. The separation of the different constituents is carried out using a DB5 capillary column (30m x 0.25 mm), (film thickness 0.25  $\mu$ m) under the following experimental conditions: carrier gas: (helium: 1 mL.min-1), energy ionization (70 eV), injector temperature (280 °C), detector temperature (280 °C). The oven is programmed at 50 °C for 5 minutes with a gradient of 5 °C.min-1 from 50 to 300 °C, 5 min at 300 °C with an injection mode split1-10.

#### **Identification of constituents**

The different constituents of essential oil have been identified on the basis of their retention indices and their mass spectra by comparison with data from the literature [48; 49; 50].

#### Hemi-synthesis of menthone 2,4-dinitrophenylhydrazone

2,4-Dinitrophenylhydrazine reagent and essential oil from the species *Mentha piperata* are used for the preparation of menthone 2,4-dinitrophenylhydrazone. The method used was reported by Leclercq [51]. It consists in dissolving 0.25 g of 2,4-dinitrophenylhydrazine in 5 mL of methanol, followed by the addition of 0.5 mL of concentrated sulfuric acid, then filtration of the lukewarm solution. To this solution is added 0.2 g of essential oil solubilized beforehand in a small volume of methanol. After about five (5) minutes, the solid formed is filtered off and

Global Journal of Pure and Applied Chemistry Research

Vol.9, No.1, pp.1-20, 2021

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

washed in a little methanol. If there is no solid, the solution is acidified with sulfuric acid. The precipitate is then recrystallized from ethanol and then dried. These conditions are summarized in Table III.

Table	III:	Operating	conditions	for	the	hemi-synthesis	of	menthone	2,4-
dinitrophenylhydrazone									

Essential oil	Quantity of reagent	Quantity of essential oil (g)	Quantity of MeOH (mL)	Quantity of H2SO4 (mL)
Mentha piperata	0.25	0.2	5	0.5

# Characterization of 2,4-dinitrophenylhydrazone Determination of melting points

The measurement of the melting temperature of the prepared hydrazone is carried out using the kofler bench. The method consists of calibrating the device with benzoic acid which has a melting point of 122.35 °C. The carriage is moved horizontally until the cursor is at the border between solid and liquid. Then the movable index is moved vertically until it indicates the melting point of the standard. The hydrazone's melting point is then taken by depositing it at the cold end of the kofler bench and moving it towards the hot zone until the first drops of liquid appear. The carriage is then moved horizontally until the cursor is at the border between solid and liquid. The moving index then indicates the melting point. Three tests are carried out,

# UV-visible spectrophotometric analysis

Analysis of 2,4-dinitrophenylhydrazone is done by a WPA Lightwawe II UV-visible spectrophotometer, connected to an HP computer.

# **Preparation of solutions**

# 2,4-Dinitrophenylhydrazine solution

The 2,4-dinitrophenylhydrazine solution is prepared by dissolving 0.3 g of 2,4dinitrophenylhydrazine in 100 mL of 0.05 M sulfuric acid solution.

#### **Essential oil solution**

The essential oil (10 to 20 mg) is placed in a 10 mL flask in which methanol is added to the mark.

#### Spectral scan

#### 2.4-dinitrophenylhydrazones can

0.5 mL of the 2,4-dinitrophenylhydrazine solution is added to 0.5 mL of the essential oil solution. An orange precipitate forms. The precipitate is left to stand for 10 minutes at room

#### Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

temperature and 5 ml of methanol are added to it: This is the solution of menthone 2, 4dinitrophenylhydrazone.

We put in the reference tank:

• 1 mL of the 30 % (V/V) water/ethanol solvent

and in the measuring tank:

• 1 mL of the 2,4-dinitrophenylhydrazone solution

#### **RESULTS AND DISCUSSION**

#### Extraction and yield of essential oils

Hydrodistillation extraction from the leaf stems of *Mentha piperata* produced the chestnut essential oil '(table IV) with a yield of 0.43 % (table V). This result is close to the results published in India [52] and Egypt [53] with respective yields of 0.42 % and 0.45 %. In addition, this yield complies with the standards described by the French Standardization Association (AFNOR) in 2000 which include the yield of essential oil of *Mentha piperata* between 0.38 % and 1.2 %.

Table IV: Organoleptic characteristics of the essentia	al oil of <i>Mentha piperata</i> L.
--	-------------------------------------

	AFNOR standards (AFNOR, 2000)			Essential		
Species	Aspect	Colour	Odour	Aspect	Colour	Odour
Mentha piperata L.	Mobile liquid	Yellow pale	Fresh characteristic more or less minty depending on the origin	Liquid	Chestnut	Minty

Table	V:	Extraction	vield	of t	he	essential	oil	of Mentha	рi	perata	L.
Labie	••	Lautuon	,1010	UI U	110	coscillia	011	01 1/10/1///	Pu	peraia	<b>_</b> .

Species	yield %						
	Our study	Previous studies/origin	Standards (AFNOR, 2000)				
Mentha piperata L.	0.43	0.42 [52]/Inde 0.45 [53]/Egypte	0.38 à 1.2				

# Physico-chemical parameter

#### **Refractive index**

The refractive index obtained at 20 °C of the essential oil of *Mentha piperata* is 1.4707 (table VI). Analysis of this table shows that this index is high. Indeed, the refractive index values of essential oils are generally high, higher than those of water at 20 °C (1.333) and olive oil at 20 °C (1.4684).

## Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

The value of this refractive index at 20 °C complies with AFNOR standards; this proves its good quality. Indeed, the standards of the French Standardization Association (AFNOR, 2000) set the refractive index of *Mentha piperata* between 1.4600 and 1.5000. Also, by comparing the value of our index to those of the literature, we find that our value is close to the values of the literature. This is the case for the refractive index value of the essential oil of *Mentha piperata* from Egypt whose IR<sub>20°C</sub> = 1.4927 [53].

## Acid index

The acid value of *Mentha piperata* essential oil is 0.672 (table VI). These values are low, less than unity. A low value of the acid number of less than 2 is proof of good preservation of the essential oil [54], that it does not deteriorate or degrade. That is, the oil is stable and is not oxidized. The essential oil stored in anti-actinic tinted bottles protected from light are protected from all attacks and possible reactions. Light, in fact, promotes the alteration of the structure of essential oils and the development of acids. Fresh essential oil is low in free acids | 54]. The value of this index compared to that of previous studies shows little difference. Indeed, the acid index of *Mentha piperata* from Egypt Ia = 0.560 [53] for which the value is also low and less than 2.

Species	Refractive	Acid	AFNOR Standards		Previous studies		
	index at 20	index	Refractive	Acid	Refractive	Acid	
	° C		index	index	index	index	
Mentha	1.4707	0.672	1.4600-	2	1.4927[53]/	0.560[53]/	
piperata			1.5000		Egypte	Egypte	
L.							

Table VI: Refractive index and acid index of essential oil of Mentha piperata L.

# Chemical composition

# Mentha piperata

The results of the analysis of the essential oil extracted from the leaf stems of *Mentha piperata* are shown in table VII. Figures 3 and 4 show the chromatograms (GC-MS) and (GC-FID) of the essential oil. Chemical analysis revealed the presence of thirty (30) constituents of which twenty (20) was identified representing a proportion of (93.26 %) of the chemical composition of the total essential oil. The essential oil consists mainly of monoterpenes (92.47 %) with a predominance of oxygenated monoterpenes (89.43 %) dominated by menthol (44.08 %), followed by menthone (14.49 %), menthyl acetate (13.54 %), mint furanone 1 (7.81 %) and menthofuran (3.04 %). Hydrocarbon monoterpenes occupy a low level of (3.04 %) characterized by limonene (2.71 %), orthocymene (0.37 %),  $\beta$ -pinene (0.35 %) and  $\alpha$ -pinene (0.15 %). Concerning the group of sesquiterpenes, this represents a minimal rate (0.79 %), oxygenated compounds are predominant (0.57%) marked by caryophyllene oxide (0.40 %) and viridiflorol (0.17 %). We note the presence of a single hydrocarbon sesquiterpene, E caryophyllene occupying a small proportion of (0.22 %). The chemical composition of this

#### Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

sample presents a qualitative resemblance to that of the essential oil of *Mentha piperata* described in Iran [55] dominated by menthol (53.28 %), menthyl acetate (15.10 %), menthofuran (11.18 %). Menthol is the major characteristic component of most of the essential oils *of Mentha piperata* in the world [56].

On the other hand, this chemical composition is qualitatively different from that of another sample of Iranian origin [57] with  $\alpha$ -terpene (19.70 %), Isomenthone (10.30%) and  $\beta$  caryiophyllene (7.60 %) as major components. Also the presence of piperitone oxide (16.00 %), 1, 8 cineole (7.03 %),  $\alpha$ -3-carene (3.27 %) in the extract of Indian origin [52] shows a different essential oil. qualitatively from our extract. Furthermore, there is also a qualitative difference with essential oil of Moroccan origin which is rich in linalool (60.72 %) and lynalyl acetate (20.79 %) [58].



Figure 3: Chromatogram (GC-MS) of essential oil of Mentha piperata L.

Global Journal of Pure and Applied Chemistry Research

Vol.9, No.1, pp.1-20, 2021



Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

Figure 4 : Chromatogramm (GC-FID) of essential oil of Mentha piparata L.



Figure 5: Chemical structures of major compounds in the essential oil of Mentha piperata L.

N°	Compounds	IK	Mentha piperata L.
1	α-Pinene	933	0.15
2	β-Pinene	977	0.35
3	Orthocymene	1026	0.37
4	Limonene	1030	2.17
5	1,8 cineole	1033	1.36
6	Cis sabinene hydrate	1069	0.34
7	Menthone	1157	14.29
8	Menthofurane	1166	3.04
9	Isomenthone	1168	2.81
10	Neomenthol	1174	0.02
11	Menthol	1180	44.08
12	Pulegone	1242	0.55
13	Piperitone	1259	0.20
14	Menthyl acetate	1293	13.54
15	Isomenthone acetate	1308	0.35
16	E Caryophyllene	1426	0.22
17	Mint furanone 1	1508	7.81
18	Mint furanone 2	1539	0.84
19	Caryophyllene oxyde	1589	0.40
20	Viridiflorol	1603	0.17
Total Oxyg Hydr Oxyg Hydr	compounds identified enated monoterpenes ocarbons monoterpenes enated sesquiterpenes ocarbon sesquiterpenes	93.26 89.43 3.04 0.57 0.22	
Aliph	atic compounds	-	

**Table VII:** Chemical composition of essential oil of Mentha piperata L.

# Hemi-synthesis of menthone 2, 4-dinitrophenylhydrazone Physical characterization

Table VIII shows the results of the hemi-synthesis of menthone 2,4-dinitrophenylhydrazone and their measured melting points. The reaction of 2,4-dinitrophenylhydrazine in the presence of sulfuric acid on the essential oil of *Menha piperata* gives rise to an orange-colored precipitate. This color is characteristic of menthone 2,4-dinitrophenylhydrazone. *Mentha piperata* essential oil contains a diverse range of chemical constituents, menthone 2,4-dinitrophenylhydrazone forms after five (5) minutes, kinetics are slow so extraction is also slow. In addition, the hemi-synthesis yield of 2,4-dinitrophenylhydrazone menthone is 29 %. This yield is low, and could be explained by the influence of the different constituents of the essential oil during the reaction. Indeed, the presence of the different constituents in the

# Global Journal of Pure and Applied Chemistry Research

Vol.9, No.1, pp.1-20, 2021

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

essential oil changes the speed of the extraction, resulting in slow kinetics. This slowing down means that there is competition or hindrance between the compound to be extracted and the other constituents of the mixture. This Competition is linked to steric and electronic effects [59]. The melting point of hydrazone in *Mentha piperata* essential oil is 145 °C. This melting point is approximate to that of the literature [60], and corresponds respectively to that of menthone 2,4-dinitrophenylhydrazone.

Table VIII: Yield and physical properties of the menthone 2,4 dinitrophenylhydrazone

Compound of essential oil converted into 2,4- dinitrophenylhydrazo- ne	Aspect	Colour	Yield (%)	Melting point Measured (°C)	Melting point of literature (°C)
<i>Mentha piperata</i> L. (menthone)	Précipitate	Orange	29	145	146

# Characterization by UV-visible spectrophotometry Menthone 2,4-dinitrophenylhydrazone

The maximum wavelength of menthone 2,4-dinitrophenylhydrazone derived from the essential oil of *Mentha piperata* of 356 nm was recorded (table IX). This value is characteristic of the C=N chromophore of 2,4-dinitrophenylhydrazones whose maximum absorption wavelength is  $(\lambda max = 360-370 \text{ nm})$  [61; 62; 63].

**Table IX:** Maximum absorption wavelength of menthone 2,4-dinitrophenylhydrazone

2,4-dinitrophenylhydrazone derivative	λmax (nm)
Menthone 2,4-dinitrophénylhydrazone	356

The UV-visible spectrum of menthone 2,4-dinitrophenylhydazone (figure 6) shows two main bands:

- A low intensity band at 270 nm corresponding to the  $n \rightarrow \sigma^*$  transition relating to the aniline aromatic system;
- ↓ A very intense band at 356 nm attributable to the n →  $\pi$  \* transition of the C=N group of the hydrazone.



Figure 6: UV-visible spectrum of menthone 2,4-dinitrophenylhydrazone

The different absorption bands of menthone 2,4-dinitrophenylhydrazone, their chromophores and their groups are shown in table X.

Table	X:	Different	absorption	bands of	menthone 2	2.4-dinitro	ophenvlhvdra	azone
I abic	<b>/\</b> •	Different	absorption	Danus OI	menuione 2	2, <b>4-</b> uiiiiii (	phenymyura	LOIIC

2,4- dinitrophénylhydrazone derivative	Absorption band λ(nm)	Transition et Chromophore	Group
Menthone 2,4-	270	n→σ* (C-NH-)	aniline aromatic
dinitrophénylhydrazone	Not very intense		system
	356	$n \rightarrow \pi^* (C=N)$	Hydrazone
	Very intense band		

# 2,4-dinitrophenylhydrazine

The UV-visible spectrum of 2,4-dinitrophenylhydrazine shown in figure 7 shows three main bands at 346 nm, 360 nm and 377 nm due to chromophoric groups C-N and NO<sub>2</sub> substituted at the aromatic ring.



Figure 7: UV-visible spectrum of 2,4-dinitrophenylhydrazine

# **CONCLUSION:**

*Mentha piperata* L. essential oil is an abundant and important source of menthone, a high added value molecule useful in the perfume, pharmaceutical, cosmetic and chemical industries. This interest has motivated our efforts to transform this molecule into hydrazone in order to enhance its economic and biological value. Menthone 2,4-dinitrophenylhydrazone was hemisynthesized by a simple, easy method from essential oil of *Mentha piperata* with a conversion rate (yield) of 29 %, in a short time of five (5) minutes. Analysis of menthone 2,4-dinitrophenylhydrazone by UV-visible spectrophotometry showed an absorption maximum wavelength of 356 nm. The UV-visible spectrophotometric method employed for the determination of this hydrazone is convenient, fast and simple. The hemi-synthesized menthone

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

2,4-dinitrophenylhydrazone could be useful in the pharmaceutical, agrochemical industries [64] and in cosmetics.

#### **REFERENCES** :

- [1]. Jahandiez E. et Maire E. (1934). Catalogue des plantes du Maroc, spermaphytes et pteridophytes. Tomes III P. Lechevalier, librairie 12, rue de towenon Vie, Alger-Paris.
- [2]. Derwich E. ; Benziane Z. ; Taouil R. ; Senhaji O. and Touziani M. (2010). Aromatic plant of Marocco: GC/MS analysis of essential oils of leaves of *Mentha piperata*. Advances in Environnemental Biology, '(1): 80-85.
- [3]. Mei-Lin Tsai; Chin-Tung Wu; Tsen-Fang Lin; Wei-Chao Lin; Yu-Chun Huang and Chao-Hsun Yang. (2013). Chemical Composition and biological properties of essential oils of two mint species. *Tropical Journal of Pharmaceutical Research*, 12(4): 577-582.
  - [4].Ndzeli Likibi B., Gouollaly Tsiba; Madielé A. B.; Nsikabaka S. Moutsamboté J.M.;
    Ouamba J. M. (2015). Constituants chimiques de l'huile essentielle de *Mentha piperata* L.
    (Lamiaceae) du Congo. *Journal of Applied Biosciences*, 92 : 8578-8585.
- [5].Guichard E. ; Genot C. ; Voilée A. Texture et flaveur des aliments : vers une conception maîtrisée. Editions 2012. Dijon, Educargi éditions, 2012, 298p.
- [6]. Haahr A. M. ; Bardou A. ; Tomsen C. E. ; Jensen S. B. (2004). Release of peppermint flavour compounds from chewing gum: effect of oral functions. *Physiology &Behavior* 82(2): 531-540.
- [7]. Vollarhd K.P.C.; Schore N. E. (1995). Traité de chimie organique, 2ème édition Bruxelles: Deboeck Université, PP: 629-674.
- [8]. Quédraogo I.W.; Boulvin M., Flammang R.; Gerbaux P. and Bonzi-Coulibaly Y. L. (2009). Conversion of natural aldehydes from *Eucalyptus citriodora*, *Cymbopogon citratus* and *Lippia multiflora* into oximes: GC-MS and FT-IR analysis. *Molecules*, 14: 3275-3285.
- [9]. Adjeroud Y. (2017). Synthèse et étude structurale d'une nouvelle famille de molécules hétérocycliques à visée médicale. Thèse de doctorat, Université Badji Mochtar Annaba, Annaba, Algérie, 159p.
- [10]. Hany S. I.; Soha R.; Abdelhadi et Haten A. A. (2015). Hydrolysis and hydrazinolysis of isatin-based-ald-and ketazines. *Journal of Chemistry*, 2015: 1-6.
- [11]. Rauk A. (2001). Orbitale interaction of organic chemistry, 2<sup>nd</sup> édition, John wiley and sons, New York.
- [12]. Greene T.W.; Wuts P.G.M. (1999). Protective groups in organic synthesis, 3<sup>rd</sup> éd. Wiley, New york, 355 pages.
- [13]. Shriner R.L.; Fuson R.C.; Curtin D.Y. & Morill T.C. (1980). The systematic identification of organic compounds, 6<sup>th</sup> éd., (John Wiley, New York).
- [14]. Armbruster F.; Klingebiel U.; Noltemeyer M. and Naturforsch (2006). 61b, 225.
- [15].Shobha B. (2011). Synthesis, caracterization and spectrophotometric determination of fer
   (II) complexe of 2,4-dihydroxybenzaldehydeisonicotinoylhydrazone(E)-N-(2,4-dihydroxybenzylidène)isonicotinohydrazide, it's application and biological activity. *Pelagia Research Library. Der Chemica Sinica*, 2 (4): 64-71.

#### Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

- [16]. Battula S.R.; Som S. D. and KIran B. (2012). Determination of molybdenium (VI) in *Arnaranthus* and *patato* by new extractive spectrophotometry method with isonitro pisopropyl acetophenone phenyl hydrazone. *Research Journal of Pharmaceutical*, *Biological and Chemical Sciences*, 3(1): 580-584.
- [17]. Jain J.; Kumar Y.; Sinha R.; Kumar R. (2011). Menthone aryl acid hydrazone: a new class of anticonvulsant. *Medicinal Chemistry*, (7(1): 56-61.
- [18]. Sammaiah G.; Srivaslava R.S. (2004). Synthesis and pharmacology evaluation of new menthone derivatives. *Indian Drugs*, 4(1(10): 624-627.
- [19]. Nesterkina M.; Barbalat D. Menthone derivatives with anticonvulsant effet. 5<sup>th</sup> *International Electronic Conference on Medicinal Chemistry*, Turkey, 2019.
- [20] Al-khattaf F.S.; Arunadevi M.; Ashraf A. H.; Idhayadhulla A. (2021). Antimicrobienne and cytotoxic acivities of isoniazid connected menthone derivatives and their investigation of clinical pathogens causing infectious disease. *Journal of Infection and Public Health*. 14(4): 533-542.
- [21]. Pandeya S. N.; Singh S. B.; Amit C. (2014). Synthesis and antifungal activity of hydrazine derivatives of some carbonyls compounds. *Shrinkhala*, Vol. II (Issue II): 11-14
- [22]. Sanchit S.; Kanishk; Chhama S.; Rajender Y. (2015). Biologically active pyrimidine hydrazones. *Asian Journal of Pharmaceutical Research*, 5(3): 162-165.
- [23]. Mohoreb R. M.; El-Sharkawy K. A.; Al Farouk; Fatma O. (2019). Synthesis, cytotocxicity against cancer and normal cell lines of novel hydrazide hydrazone derivative bearing 5Hchromeen-5-one. *Medicinal Chemistry Research*, 28: 1885-1900.
- [24]. Mishra V.; Pandaya S. N.; Pannecouque; Witvrouw M.; Declerq E. (2002). Anti-HIV activity of thiosemicarbazone and semicarbazone derivatives of (±) -3-menthone. *Chemistry in Life Sciences*, 335(5):183-186.
- [25]. Mishra V. Pandaya S.N. (2001). Analgesic activity and hypnosis potentiation effect of (±)
   -3-menthone semicarbazone and thiosemicarbazones derivatives. *Acta Pharm.*, 51: 83-88.
- [26]. Ansari A.. Manssouri M.; Znini M.; Lakbaiki Z.; Azrour M. (2020). Experimental theoretical study on corrosion inhibition of new synthesized (menthone derivative/menthopyrazole compounds) for mild steel in 1M HCl solution. *Mediterranean Journal of Chemistry*, 10(1):62-76.
- [27]. Zalaru C. M.; Zalaru F.; Ceabanu A.; Mairinescu M; Raiciu A. D. (2020). Coordination compounds of cu (II) and Ni(II) with schiff bases derived from formylmenthone and aromatic amines. Preprints (www. Preprints. Org).
- [28]. Nesterkina M.; Barbalat D.; Zheltvay I; Rakipov I. (2019). (2S, 5R)-2-isopropyl-5methylcyclohexanone hydrazones. *Molbank. Communication*. p 1-6.
- [29]. Mullica D. F.; Garner C.M.; Prince M. E. Prince; Mossman B.C. and Sappenfield E. I. (1992). Structure analyses of menthone and isomenthone tosylhydrazone, C<sub>17</sub>H<sub>26</sub>N<sub>2</sub>O<sub>2</sub>S. *Journal of Crystallographic and Spectroscopic Research*, 22(5): 515-518.
- [30]. Srivastav S.; Shukla C.; Luhash K.; Pandeya S. N. (2016).Pyridimine hydrazone as a potent anticonvulsant. *World Journal of Pharmaceutical Research*, 5(6): 2101-2108.

- [31]. Yang D., Luo X. (2012). NMR characterization of reaction product of L-menthone hydrazine. *Chinese Journal of Magnetic resonance*, vol. 29: 401-409.
- [32]. Srivastav S.; Mindhaj M. A.; Probhakar M.K.; Luhach K. (2021). Synthese of potential & biologically active pyridimidine hydrazones derivatives. *World Journal of Pharmaceutical Research*, 4(10):959-967.
- [33]. Sakuragawa A.; Yoneno T.; Inoue K. and Okutani T.(1999). Trace analysis of carbonyl compounds by liquid chromatography-spectrometry after collect as 2,4-dinitrophénylhydrazone derivatives. *Journal of Chromatography A.*, 844(1-3):403-408.
- [34]. Zweiner C.; Glaumer T.and Frimmel F.H (1997). Method optimization for the measurement of carbonl compound in disinfected water by 2,4-dinitrophenylhydrazine derivation and LC-ESI/MS/MS. *Analytical and Bioanalytical Chemistry*, 372(5-6): 615-621.
- [35]. Soraya de M. Ochs; Maira Fasciotti and Annibal D. P. N. (2015). Hydrazones of carbonyl compounds by RRLC-UV and RRLC-MS(/MS): A comparison of methods. *Journal of Spectroscopy*, 2015, 11 pages.
- [36]. Yasuhara A.; Tanaka Y.; Makishima M.; Suzuki S. andShibamoto T. (2011). LC-MS analysis of low molecular weight carbonyl compound as 2,4-dinitrophenylhydrazones using negative ion mode electrons pray ionization mass spectrometry. *Journal of Chromatography Separation Technique*, 2(1): 1-5.
- [37]. JI-Zhou D. and Serban M. (2004). Gas chromatography-mass spectrometry of carbonyl compounds in cigarette mainstream smoke after derivatization with 2,4 dinitrophenylhydrazine. *Journal of Chromatography A* 1027(1-2):25-35.
- [38]. Dai Tian-You, Wang Qin-hui; Wei Fu-Sheng (1999). Sampling and determination of carbonyl in stack gas. *Journal of Environmental Science*, 11(1): 48-56.
- [39]. Suze M. V.L.; Laurens H. and Uwe K. (2004). Determination of aldehydes and ketones Using derivatization with 2,4-dinitrophenylhydrazine and liquid chromatographyatmospheric-pressure photoionization-mass spectrometry, *Journal of Chromatography A*, 1058 (1-2): 107-112.
- [40]. Gori G.; Meneghetti P.; Sturaro A;; Parvoli G.; Doretti L. & Bartolucci G.B. (1995). High performance liquid Chromatographic determination of methyl ketones in urine as its methyl-2-benzothiazolinone derivative. *Chromatographia*, 40: 336-340.
- [41]. Myadagbadam U.; Meng G. D.; Wang Y.; Hasi S.; Edernechimeg C.H.; Chimedragchaa C.H. 2019. HPLC method for determination and optimization of 2,4-DNPH Pre-culumn derivatization effect of reaction conditions for muscone in natural musk. *Natural Products Chemistry & Research*, Vol. 7, Iss. 2, No: 364.
- [42]. Stenlake J.P. and Williams W. D. (1957). The spectrophotometric determination of  $\alpha$ , $\beta$ unsaturated aldehydes and ketones with Girard-T-reagent. *Journal of Pharmacy and Pharmacology*, 9(1): 900-907.
- [43].Praven P-sai; Anupam B.; Jagathi V.; Devala Rao G. (2010). Spectrophotométric determination of tolperisone using 2,4-dinitrophénylhydrazine reagent. *Int. J. Res. Pharm. Sci.*, 1(3): 317-320.

- [44]. Disha N. S. and Gurupadhayya B. M.(2015). Spectrophotometric determination of Sitagliptin phosphatein bulk and pharmaceutical formulations. *International Journal of Innovative Science, Engineering & Technology*, 2(7): 702-709.
- [45]. Clevenger J. F. (1928). Apparatus for the determination of volatile oil. *Journal of the American Pharmaceutical Association*, 17(4): 45-349.
- [46]. AFNOR : Association Française de Normalisation. (2000). « Recueils des normes : les huiles essentielles. Tomes 2. Monographies relatives aux huiles essentielles». AFNOR, Paris, 661-663.
- [47]. AOAC.(1990). Official methods of analysis (15 th ed.) Washington., VA. Association of Official Analytical Chemists.
- [48]. Adams R. P. (2001). Identification of essential oils by gas chromatography quadrupole mass spectroscopy. Carol Stream. IL, USA: Allured Publishing Corporation. 101p.
- [49]. Joulain D.& Kőning W. A. (1998). The atlas of spectral data of sesquiterpene hydrocarbons. Hamburg Germany: EB-Verlag.
- [50]. Davies NW. (1990). Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and carbo wax 20M phases. *Journal of Chromatog.*, 503: 1.
- [51]. Leclercq (2008). Détermination de la teneur en composés toxiques dégagés lors de la combustion d'encens et des bougies parfumées. Rapport du projet d'étude. Université de Rouen, *Institut National des Sciences Appliquées*. 52p.
- [52]. Sharma V.; Sharma N.; Pathania V.; Malika R.; Singh B.; Guptac C. R. (2010). GC-MS analysis and anti-microbial activity of essential oil of *Mentha piperata* L. from Kulla-A north India region of higher altitude Himalayas. *International Journal of Drug Development and Research*. 4:134-140.
- [53]. Ayaidia B. (2011). Etude comparative de tris variétés d'huiles essentielles de Menthe dans la region Ourgla. Mémoire de fin d'étude, Université Kasdi Marbah, Ourgla, Egypte. 48p.
- [54]. Fauconnier M. L. (2006). HE d'Ylang Ylang: sa fiche qualitative et son suivi –exposé pour les GIE maison des Epices des Comores
- [55]. Mohammad J.S., Marjan M.,Kamiar Z.,Keyvan P., Ramin M. and Kimia H.(2012). Chemical composition, antitifungal and antibiofilm activity of the essential oil of *Mentha piperata* L. *International Scholarly Research Network Pharmaceutics*. P.1-6.
- [56]. Foster S.(1996). Peppermint: *Mentha piperata*. *American Botanical Council-Botantical Series*, 306: 3-8.
- [57]. Rasooli I.; gachkar L., yadegarinia D., Bagher R. M.; Alipoor A. S. D. (2008). Antibacterial and antioxidative characterization of essential oils from *Mentha piperata* and *Mentha spicata* grown in Iran. *Acta Alimentaria*, 38: 41-52.
- [58]. Debbab A., Mosaddak M. (2007). Chemical characterization and toxicological evaluation of the essential oil of *Mentha piperata* L. growing in Morocco. Studii Si Cercetari Stiintifice: *Chimie si Ingenirie Chimiaca, Biotech., Ind. Alim.* 8(3): 281-288.
- [59]. Kasséhin C. U.; Gbaguidi F.A.; Kapanda G.N.; McCurdy C.; Bigot A. K. and Poupaert J. H. (2013). Electronic and steric effects in the control of anilinium chloride catalyzed

Print ISSN: ISSN 2055-0073(Print), Online ISSN: ISSN 2055-0081(Online)

condensation reaction between aldones and 4-phenylthiosemicarbazide. *African Journal of Pure and applied chemistry*, 7(9):325-329.

- [60]. Rappoport Z. (1967). CRC Hand book of tables for organic compound identification 3<sup>rd</sup> edition P.147.
- [61]. Zhou X., Mopper K.(1990). Measurement of sub-part-per-bilion levels of carbonyl compounds in marine air by a simple cartridge trapping procedure followed by liquid chromatography. *Environmental Science and Technology*; 24: 1482-1485.
- [62]. Pötter W., Lamotte S.; Engelhardt H. karst U. (1997) Non-porous silica ultrafast reversed phase high performance liquide chromatography separation of aldehydes and ketones 2,4-dinitrophénylhydrazones. *Journal of Chromatography* A. 786: 47-55.
- [63]. Levart A., Veber M. (2001). Determination of aldehydes and ketones in air samples using criptrapping sampling. *Chemosphene*; 44:701-708.
- [64]. Anwar T.M. AL-Thib and A. Khudhair (2016). Spectroscopic study of some aromatic hydrazones derivated from substituted benzophenones and benzaldehydes. *PCAIJ*, 11(1): 024-034.