
Sensitivity Analysis and Future Farm Size Projection of Bio-Fortified Cassava Production in Oyo State, Nigeria

¹Kolapo Adetomiwa* and ²Olayinka. J. Yusuf

¹Department of Agricultural Economics, Faculty of Agriculture,
Obafemi Awolowo University, Ile Ife, Osun State, Nigeria.

²Department of Agricultural Economics and Extension Services, Kwara State University, Malete,
Nigeria.

Citation: Adetomiwa K and Yusuf O.J. (2022) Sensitivity Analysis and Future Farm Size Projection of Bio-Fortified Cassava Production in Oyo State, Nigeria, *European Journal of Agriculture and Forestry Research*, Vol.10, No.3, pp. 35-50

ABSTRACT: *The study examined the costs and returns to bio-fortified cassava production and forecast the future farm size of bio-fortified cassava production in the study area. A multistage sampling technique was used to select 150 respondents in the study area. Primary data were used for the study which were collected through a well-structured questionnaire. Data collected were analyzed using descriptive, Markov chain, and budgetary analysis. The result of the study showed that the mean age of the respondents were 47(±13.77) with a mean years of experience of 14.62(±6.92). the result of the study showed that TMS 01/0593, TMS 01/0539 and TMS 01/0220 were the mostly grown varieties of bio-fortified cassava varieties in the study area. The result of the budgetary analysis showed that the average net return (net farm income) from the production of bio-fortified cassava was ₦196710.95 with RORI of 224.95%. The result revealed that at 35% increase in cost of production, the rate of return on investment dropped to 140.70% in which the investment will not be viable. The bio-fortified cassava farmers have a great potential to boost production through increases in farm sizes of the bio-fortified cassava famers until the year 2026 when equilibrium would be attained at about 2.85ha. in other to adequately achieve these goals, more improved varieties of bio-fortified cassava should be provided, and also, infrastructures should be put in place to help boost farmers moral in their cause of production in the study area.*

KEY WORDS: sensitivity analysis, bio-fortified cassava, Markov chain, farm size, Oyo State.

INTRODUCTION

Cassava is an important staple food in Nigeria. Cassava is a starchy crop which contributes to the staples of millions in sub-Saharan Africa (SSA). According to Otekunrin and Sawicka (2019), about 177,948 million tonnes of cassava were produced in Africa. Nigeria is regarded as the world's largest producer of cassava with a total of about 20.4 percent of the world export in year 2017 (Otekunrin and Sawicka, 2019). Cassava is a major staple food crop in Nigeria. As defined by Otekunrin and Sawicka (2019), a staple crop is the one that is been eaten regularly and which also provides larger proportions of the population's nutrients. Cassava fulfil this purpose as it can be eaten raw or in a processed form. Cassava is an essential component of the diet of about 70

million Nigerians (FAO, 2013). Nigeria, being the largest producer of cassava in the world is producing an average annual estimate of 45 million metric tons which had been translated into a major global market share of about 19 percent (Hillocks, (2002); Phillips *et al.*, 2004). The production of biofortified vitamin-A cassava started in 2011 with the intervention of the International Center for Tropical Agriculture (CIAT) and the International Institute of Tropical Agriculture (IITA) which were funded by Harvest Plus program. Five years after the intervention program, statistics revealed that over 1million of Nigerian farming households grows yellow cassava varieties that contains substantial quantities of vitamin-A even after processing. In Nigeria diets today, yellow bio-fortified cassava represents additional source of vitamin A (Saltzman *et al.*, 2014).

However, majority of the bio-fortified cassava farmers were still producing on a small scale due to a myriad of factors including lack of general acceptance of the product despite its nutritional benefits to human population. According to Ilona *et al.*, (2017), production of bio-fortified cassava is still on a relatively small farm size despite the release of the first wave of vitamin A cassava in 2011. Hence, to a large extent accounts for the low supply of bio-fortified cassava products like vitamin A yellow garri, fufu and high-quality cassava flour relative to the demand for these products thus necessitating the need to project the future farm size with a view to determining what the future holds for bio-fortified cassava production in the study area.

In commercial enterprises, profit is a major motivating factor. The profit and profitability levels of farm enterprises may be influenced by the farm size. This is because it is assumed that with larger farm size the cost of production is spread across the number of hectares and as such profitability is increased. Thus, it is necessary to examine the trend in size of farms to determine their intertemporal performance. The specific objective of the study were to described the socio-economic characteristics of the bio-fortified cassava farmers in the study area; forecast the future farm size of bio-fortified cassava production and estimate the cost and return to bio-fortified cassava production in the study area.

Theoretical framework

In other to project the future farm size of bio-fortified cassava production, Markov chain model was utilized. Markov chain are one of the conceptual devices used in analyzing the types of changes obtainable when there is movement from one state to another (Anders, 2016). It was first used in the study of Markov A.A in 1907 and has been used in various sectors ranging from agriculture, health and migration studies to forecast and predict future trend. In Agriculture it is useful in predicting and forecasting the behavior of farmers as they move from one categories of farm size to another. It is one of stochastic process in which the probability or likelihood associated with a set of possible future outcome is stated. A stochastic process refers to mathematical model with a sequence of random variables which assumes that any population of individuals or firms can be classified into various groups. As such, movements between states over time is regarded as a stochastic process (Olatidoye *et al.*, 2018). A finite Markov process is one in which the outcome of a given trial (experiment) in the time (t + 1) essentially depends on the outcome of the trial in the preceding time period (t) and this dependence holds at all the various stages of the trial.

Markov chains are often characterized by the dynamic property, such that as the present condition is known, prediction about the future outlook or behavior of the process remain the same, even if additional information about past history of the process is known (Anders, 2016). Finite Markov chain process often determined by specification of a given set of states ($S_1, S_2 \dots S_n$). Only one state is achievable at a given time and it moves progressively from one state to another. The probability of moving from S_i to S_j is given for every pair category can be represented in the form of transition matrix P . P_{ij} refers to the probability of moving from S_i to S_j in the next step. The element of the matrix must be non-negative, and the row sum of the elements is one. when all the initial probability is known, outcome of the n th step, can be gotten.

$$P = \begin{pmatrix} & S_1 & S_2 & \dots & S_n \\ S_1 & P_{11} & P_{12} & \dots & P_{1n} \\ S_2 & P_{21} & P_{22} & \dots & P_{2n} \\ \vdots & \vdots & \vdots & & \vdots \\ S_n & P_{n1} & P_{n2} & \dots & P_{nn} \end{pmatrix} \dots\dots\dots (1)$$

$$P = [P_{ij}] = \left[\frac{n_{ij}}{\sum_{j=1}^n n_{ij}} \right] \geq 0 \quad (j = 1, 2, 3, \dots, m) \dots\dots\dots(2)$$

Hence, the future path of the stochastic process is given by;

$$\begin{aligned} P^{(0)} P &= P^{(1)} \text{ state vector in time, } t + 1 \\ P^{(1)} P &= P^{(2)} \text{ state vector in time } t + 2 \\ P^{(m-1)} P &= P^{(m)} \text{ state vector in time } t+m \dots\dots\dots (3) \end{aligned}$$

Therefore, $P^{(0)} P^{(e)}$ gives the fixed probability vector, or equilibrium probability vector of the stochastic process.

$$\begin{aligned} \text{Hence: } P^{(m)} &\rightarrow P^{(e)} \text{ as } m \rightarrow \infty \\ P^{(0)} P^{(e)} &= P^{(e)} \\ P^{(e)} P &= P^{(e)} \dots\dots\dots (4) \end{aligned}$$

The equilibrium farm size indicates that the number of people entry a particular category of farm size is equal to the number of farmers leaving the group. The underlying assumptions on which Markov chain includes the following; The structure of the population when the transition

probability is made remain constant, the underlying determinant of a change in one category of farm size is represented by a probability of individual movement from one category of farm size to another depends on the result proceeding of the period.

There are several application of Markov model in Agricultural economics such as market structure and economic development. Empirical studies that employed the use of Markov chain include those of (Alimi *et al.*, 2007; Baruwa *et al.*, 2011; Olatido *et al.*, 2018).

METHODOLOGY

Area of Study

The study was conducted in Oyo States, Nigeria. Oyo State is an inland state in South-Western Nigeria, with its capital at Ibadan. It is bounded in the north by Kwara State, in the East by Osun State, in the South by Ogun State and in the West partly Ogun State and partly by the Republic of Benin with a population of 5,591,589 people (NPC, 2006). Oyo State is homogeneous, mainly inhabited by the Yoruba ethnic group who are primarily agrarian but have a predilection for living in high-density urban centers. Oyo State covers approximately an area of 28,454 square kilometers. Oyo State is located in the rainforest vegetation belt of Nigeria on longitude of $2^{\circ}38.66^{\circ}\text{N}$ and $4^{\circ}38.25^{\circ}\text{N}$ and latitude $9^{\circ}8.74^{\circ}\text{E}$ and $7^{\circ}1.68^{\circ}\text{E}$. Agricultural activities in Oyo State include the production of different varieties of arable food crops since the climatic conditions support the production of various food crops including cassava, maize, groundnut etc. A large proportion of the bio-fortified cassava were being produced in the State as the distribution of bio-fortified cassava stem started in Oyo State in 2011, hence the choice of the study area.

Sampling procedures and sample size

Multistage sampling procedures were employed for the study. The first stage involved purposive selection of two Local Government Areas (LGAs) because of the concentration of bio-fortified cassava producers in the areas. The second stage involved random selection of three communities from each of the selected LGAs. At the third stage, twenty-five bio-fortified cassava farmers were purposively selected from each community to make a total of 150 (One hundred and fifty) respondents. Primary data were used for the study. The primary data were sourced from cross-sectional survey of bio-fortified cassava farmers in the study area with the aid of well-structured questionnaire to cover information about the socioeconomic characteristics of respondent and inputs and outputs of bio-fortified cassava production. Data were collected in December 2018.

Analytical techniques

The data were analyzed using descriptive statistics, Markov Chain Analysis, Farm budgeting analysis (Gross margin sensitivity).

Descriptive statistics

Descriptive statistics was used to analyze the socio-economic characteristics of the bio-fortified cassava farmers.

Markov Chain analysis

Markov Chain analysis was employed in this study to predict and forecast the future farm size of bio-fortified cassava farmers production. Markov Chain process is a stochastic model used in the analysis of economic variable with the availability of a time-ordered data. (Amina and Akhigbe, 2017). Bio-fortified cassava farmers were grouped according to some criteria of farm sizes (states). Secondly the evolution of bio-fortified cassava farmers through these states can be regarded as a stochastic process. The probability of moving from one state (t) to another (t+1) is a function only of the two states (t, t+1) involved. The movement of the bio-fortified cassava farmers within the farm size depends on the initial farm size attained and the number of years involved which is independent of the previous history (Ander, 2016). Within this framework, farm sizes cultivated are the variable whose movement over time is to be analyzed, and the following class intervals will be used in defining the admissible states.

Table 1: distribution by size

Class	Farm size in (ha)
S1	1-2
S2	2.1-3
S3	3.1-4
S4	4.1-5

Therefore, in a year it is possible for a bio-fortified cassava farmer to be in any one of the four specified positions. Having defined the data and the ranges for each class, the year-to-year history of each bio-fortified cassava farmer in terms of his movement among the various classes was used in developing the transition matrix, which reflects the behaviour of the sample of bio-fortified cassava farmers. Let u_{ij} represent the number of farmers moving from class i to class j through the years under consideration. The transition probabilities (P_{ij}) can be represented in the form of transition matrix P . P_{ij} is the probability of bio-fortified cassava farmers transitioning from state i to j (one farm size category to the other).

$$P_{ij} = \begin{matrix} S_1 \\ S_2 \\ S_n \end{matrix} \begin{bmatrix} S_1 & S_2 & \dots & S_n \\ P_{11} & P_{12} & & P_{1n} \\ P_{21} & P_{22} & & P_{2n} \\ P_{n1} & P_{n2} & & P_{nn} \end{bmatrix}$$

$$P = [P_{ij}] = \left[\frac{n_{ij}}{\sum_{j=1}^n n_{ij}} \right] \geq 0$$

- i. $\sum_{j=1}^m P_{ij} = 1$ (row summation of probability should equal to one)
- ii. $P_{ij} \geq 0$ (for all i and j)
- iii. $P_j = \frac{n_j}{N}$ ($1 \leq j = 1, 2, 3, \dots, m$)

The long run equilibrium is attained when the total number of bio-fortified cassava farmers entering a given farm category equals the number of farmers exiting. This is expressed as follows:
 $eP = e$.

$$(e_1, e_2, e_3, \dots) \begin{bmatrix} P_{11} & P_{12} & P_{13} \\ P_{21} & P_{22} & P_{23} \\ P_{31} & P_{32} & P_{33} \end{bmatrix} = (e_1, e_2, e_3, \dots)$$

Table 2: First-order Markov model for farm size transitions

Period 1 (t)	Period 2 (t + 1)				Total
	S ₁	S ₂	S ₃S _n	
S ₁	n ₁₁	n ₁₂	n ₁₃	...n _{1m}	n ₁
S ₂	n ₂₁	n ₂₂	n ₂₃	n _{2m}	n ₂
S ₃	n ₃₁	n ₃₂	n ₃₃	n _{3m}	n ₃
.
.
.
S _m	n _{m1}	n _{m2}	n _{m3}	n _{mm}	n _m
Total (Period t + 1)	n ₁	n ₂	n ₃	n.m	N

Source: Author's, 2019

Budgetary technique (Gross margin sensitivity)

The gross margin of the farm is a measure of output and farm profitability, which is a useful indicator in planning. Gross Margin (GM) is the difference between total revenue and total variable cost while Gross Margin Sensitivity (GMS) is the difference between total revenue and changes in total variable cost. Since parameters and the output have different measurement units, they are not directly comparable. This problem can be overcome by calculating the "elasticity" or the percentage change in output to a percentage change in other parameters (Pannell, 1997). The sensitivity is calculated to explore the impact of assumptions regarding the changes in farm sizes

on the gross margin, by using the principle “what if” (Dachin *et al.*, 2016). The sensitivity is interpreted as the elasticity of gross margin to changes in farm sizes by +/- 5%, 10%, 15 and 20%).

$$GM = TR - TVC \dots\dots\dots (5)$$

$$NI = GM - TFC \dots\dots\dots (6)$$

$$ROI = NFI / TC \dots\dots\dots (7)$$

$$BCR = TR / TC \dots\dots\dots (8)$$

TVC = Summation of all the variable cost which includes;

- i. Land preparation
- ii. Planting materials
- iii. Chemical used
- iv. Labour used (planting, weeding, fertilizer and pesticide application and harvesting)
- v. Transportation

Where:

GM = Gross margin

NFI = Net farm income

TC = Total cost incurred

ROI = Return on investment

BCR = Benefit cost ratio

TVC = Total variable cost incurred

TFC = Total fixed cost incurred

TR = Total revenue generated from production

RESULTS AND DISCUSSION

Socio-economic characteristics of respondents

Presented in Table 2 were the socio-economic characteristics of the bio-fortified cassava farmers in the study area. The mean age of the respondents were 47(±13.77) which implies that bio-fortified cassava farmers in the study area were young and active thus expected to be productive. They are also expected to be open to adoption of new innovation in agricultural practices. Majority (53%) of the respondents were women. This agree with Oparinde *et al.* (2014) that cassava production in Nigeria were mostly common among the women gender. Majority (79%) of the respondents were married implying that they were responsible. It might be due to the fact that marriage is cherished in the study area. The mean years of formal education was 14.39 (±6.83) which implies that respondents in the study area were literate and thus, can read and write. The mean household size were 5.31 (±2.26). This implies that the respondents invariably had a medium

to large family size and the use of family labor is possible in the study area. About 58% of the respondents had access to credit to facilitate their production of bio-fortified cassava in the study area. These might be due to the fact that the respondents belong to association in the study area. The mean years of experience was 14.62(\pm 6.92). This implies that respondents in the study area had been into cassava production for a long time even before the introduction of new improved bio-fortified cassava in 2011. Majority (86%) of the respondents belong to one association or the other. They can thus experience group benefits such as credit facility, inputs etc.

Table 2: Socio-economic Characteristics of Bio-fortified Vit-A Cassava Farmers

Variables	Bio-fortified cassava farmers
Age (years)	47(\pm 13.77)
Female (%)	53.00
Married (%)	79.00
Formal education (years)	14.39 (\pm 6.83)
Household size (#)	5.31 (\pm 2.26)
Access to credit (%)	58.00
Years of experience (years)	14.62(\pm 6.92)
Membership of association (%)	86.00

Source: Data Analysis, 2018

Farm specific characteristics

Presented in Table 2 is the farm size of bio-fortified cassava farms in the study area the result shows that majority (41.33% and 38% for 2017 and 2018, respectively) of the respondents cultivated between 1-2ha in the two years respectively. The mean farm size in 2017 and 2018 was 2.1ha and 2.19ha. The results imply that bio-fortified cassava farms in the study area were small scale. From Table 4, about 39.33% of the respondents inherited their farm land, 22% purchased their farm land, 24% rented their farm land while 14.67% of the respondents gotten their farm land through communal/gift. About 21.33%, 25.33% and 20.66% of the respondents grown TMS 01/0593, TMS 01/0539 and TMS 01/0220 respectively. These varieties of bio-fortified cassava were the second wave of the bio-fortified cassava distributed in 2016 and was observed to be an improved variety over the ones that were first released in 2011 as listed in Table 4, hence the reason behind cassava farmers adopting this varieties in the study area. Majority (55.33%) of the respondents practiced intercropping in the study area. This implies that they tend to maximize land resources for the production of bio-fortified cassava as crops like maize were observed to be intercropped with Bio-fortified cassava in the study area.

TABLE 3: Farm size of bio-fortified cassava farms

Farm size (ha) categories	2017		2018	
	Frequency	(%)	Frequency	(%)
1-2	62	41.33	57	38.00
2.1-3	49	32.67	52	34.67
3.1-4	36	24.00	37	24.67
4.1-5	3	2.00	4	2.66
Mean	2.14		2.19	

Source: Data analysis, 2018

TABLE 4: Farm specific characteristics

Variables	Frequency	Percentage
Mode of land acquisition		
Inherited	59	39.33
Purchase	33	22.00
Rent	36	24.00
Communal/Gift	22	14.67
Varieties grown		
TMS 01/1371	16	10.67
TMS 01/1412	19	12.67
TMS 01/1368	14	9.33
TMS 01/0593	32	21.33
TMS 01/0539	38	25.33
TMS 01/0220	31	20.67
Agricultural system practiced		
Sole cropping	67	44.67
Inter cropping	83	55.33

Source: Data analysis, 2018

Markov chain analysis for bio-fortified cassava farm size

The movement of bio-fortified cassava farmers from one farm size category to another between the two periods (2017 and 2018) were presented in Table 5. The farm size was categorized into four groups; 1-2ha, 2.1-3ha, 3.1-4ha and 4.1-5ha. From Table 5, the first cell on the first row (S_1S_1) contains the number of bio-fortified cassava farmers (53) that cultivated between 1-2ha in the first period (2017) and still remained in the same category in the second period (2018). The figure in the second cell of first row (S_1S_2) represents the number of bio-fortified cassava farmers (6) in the farm size category 1-2ha in the first period but had moved to 2.1-3ha farm size category in the second period. The figure (0) in the third cell of the first row (S_1S_3) implies that no farmer in the 1-2ha farm size category in 2017 had moved to 3.1-4ha in the second period (2018). This is applicable to fourth cell and for other rows of the transition matrix (Table 5).

The transition probability matrix corresponding to the transition matrix of Table 5 is shown in Table 6. The entries in the cells on the principal diagonal of Table 6 indicate the tendency for the farmers to remain within a given category of farm size. These entries show that there was a strong tendency (0.90, 0.85, 0.89 and 0.67) for those farmers cultivating farm size (1-2ha, 2.1-3ha, 3.1-4ha and 4.1-5ha) respectively to remain there. This implies that for a proportion of 0.90 in the first cell of the principal diagonal (S_1S_1) for example, as many as 90% of the farmers remained in that category in the second period (2018). The proportion in the second cell of the principal diagonal (S_1S_2) corresponding to farmers cultivating 2.1-3ha is 0.85 which implies that 85% of the farmers that cultivated 2.1-3ha stands in 2017 remained in this category in 2018. However, the proportions in the cells to the right of each of the cells in the principal diagonal indicate the chances of moving to higher categories than that of the principal diagonal cell. Similarly, the proportions in the cell to the left of each of the cells on the principal diagonal indicate the chances of moving to lower categories than that of the principal diagonal cell. For example, the cells to the right of the first cell on the principal diagonal (S_1S_2 , S_1S_3 and S_1S_4) contain 0.10, 0.00 and 0.00 for 2.1-3ha, 3.1-4ha and 4.1-5ha respectively. This implies that the probability of farmers who cultivate 1-2ha category in period one to move to 2.1-3ha category and higher ones in period two is low.

TABLE 5: Transition matrix for farm size categories

Farm size categories 2017	2018				Total 2008
	S_1 1-2	S_2 2.1-3	S_3 3.1-4	S_4 4.1-5	
S_1 1-2	53	6	0	0	59
S_2 2.1-3	4	44	4	0	52
S_3 3.1-4	0	2	32	2	36
S_4 4.1-5	0	0	1	2	3
Total 2018	57	52	37	4	150

Source: Data analysis, 2018

TABLE 6: Transition probability matrix for farm size

Farm size categories 2017	2018			
	S_1 1-2	S_2 2.1-3	S_3 3.1-4	S_4 4.1-5
S_1 1-2	0.90	0.10	0.00	0.00
S_2 2.1-3	0.07	0.85	0.07	0.00
S_3 3.1-4	0.00	0.06	0.89	0.06
S_4 4.1-5	0.00	0.00	0.33	0.67

Source: Data analysis, 2018

Equilibrium values, actual and projected pattern of changes in farm size of bio-fortified cassava farmers

The result of the actual and projected farm size for bio-fortified cassava farmers were presented in Table 7. The projection of the structure in which the farm size of the bio-fortified cassava farmers would attain assuming the trend observed on the field during 2017 and 2018 continues over time,

implies that equilibrium will be attained in year 2026. In comparing the proportion of bio-fortified cassava farmers in different farm size in initial year with equilibrium year, the proportion of farmers in farm size 1-2ha and 2.1-3ha will decline from 0.09 and 0.70 to 0.06 and 0.30 respectively. Considering the proportion of bio-fortified cassava farmers in the farm size group of 3.1-4ha and 4.1-5ha, it would grow from 0.14 and 0.07 to 0.38 and 0.26 respectively. Furthermore, the mean farm size of bio-fortified cassava farmers on Table 7 shows an upward trend over time. At equilibrium, the mean farm size was 2.85 compared to 2.20 at the initial year in 2017. The result of this study implied that bio-fortified cassava farmers in the study area were small scale farmers but they could increase their production in the future if certain measures were put in place.

TABLE 7: Actual and projected structure of farm size among bio-fortified cassava farmers

Farm Size	2017*	2018**	2019	2020	2021	2022	2023	2024	2025	2026***	2027
1-2	0.09	0.08	0.08	0.07	0.07	0.07	0.07	0.06	0.06	0.06	0.06
2.1-3	0.70	0.65	0.59	0.51	0.48	0.45	0.42	0.33	0.30	0.30	0.30
3.1-4	0.14	0.19	0.24	0.33	0.35	0.36	0.37	0.37	0.38	0.38	0.38
4.1-5	0.07	0.08	0.09	0.09	0.10	0.12	0.14	0.24	0.26	0.26	0.26
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Mean	2.20	2.25	2.30	2.40	2.45	2.55	2.65	2.70	2.75	2.85	2.85

Source: Data analysis, 2018

*Actual year

**Starting (initial) state probability vector

***Equilibrium probability vector

TABLE 8: Average costs and return to bio-fortified cassava production per season

Variables	Amount (₦)	%of total costs
A. Total revenue	284159.27	
Variable cost		
Land preparation	42786.32	48.93
Planting material	8620	9.85
Fertilizer	1650	
Herbicides	3400	
Labor cost	14370	16.43
B. Total Variable Costs (TVC)	70826.32	81.00

Fixed cost		
Rent on land	14850	16.97
Depreciation on sprayer	843	
Depreciation on wheelbarrow	929	
C. Total fixed costs (TFC)	16622	19.00
D. Total costs (B+C)	87448.32	
E. Gross margin (A-B)	213332.95	
F. Net Farm Income (A-D)	196710.95	
Return on Investment (ROI)	2.25	
Benefit Cost Ratio	3.25	

Source: Data analysis, 2018

Costs and returns to bio-fortified cassava production

In order to ascertain the profitability of bio-fortified cassava production, the average gross margin, net returns, rate of returns and benefit cost ratio of the bio-fortified cassava farmers were calculated. The input used, costs, output data generated from the bio-fortified cassava farmers were used to compute the gross margin and net returns to bio-fortified cassava production.

The average costs and returns for the bio-fortified cassava production were presented in Table 8. The result revealed the revenue generated for one production season was ₦284159.27. From Table 8, the cost of land preparation (₦42786.32) on individual cost accounted for a large proportion (48.93%) of the total costs with the total variable costs (₦70826.32) accounting for the largest proportion (81%) of the total costs. Rent on land (₦14850) accounted for a significant proportion 16.97% of the fixed cost with the total fixed costs accounting for just 19%. The negligible small proportion of the fixed costs shows the crude method of agricultural small-scale practices in the study area. The average net return (net farm income) from the production of bio-fortified cassava in Table 8 was ₦196710.95. This implies that the production of bio-fortified cassava in the study area is a profitable enterprise.

The return on investment, indicated that for every one naira invested in bio-fortified cassava production, the farmer gains ₦2.25. The implication is that bio-fortified cassava production in the study area is profitable. The result agrees with Ogunleye *et al.* (2019) in the Profitability of investment and farm level efficiency among groups of Vitamin A cassava farmers in Oyo State Nigeria who found out that bio-fortified cassava production is a profitable business enterprise. The benefit cost ratio of 3.25 shows that for every ₦3.00 return to bio-fortified cassava production, 25k is been spent on the cost of producing the bio-fortified cassava in the study area.

Rate of Return on Investment

$$RORI = \frac{TR - TC \times 100}{TC}$$

$$= \frac{284159.27 - 87448.32}{87448.32} \times 100$$

$$RORI = 224.95\%$$

Sensitivity Analysis and Rate of Return on Investment

The rate of returns on investment of bio-fortified cassava production showed a high returns in the enterprise (224.95%). The rate of returns on investment of bio-fortified cassava production was subjected to a sensitivity analysis to establish the point at which profitability might not be certain. With respect to input, increasing the costs from +10 to +30% did not significantly impact the rate of return on investment (Table 9). Furthermore, the result revealed that at 35% increase in cost of production, the rate of return on investment dropped to 140.70%. This implies that with outmost concern, bio-fortified farmers should try as much as possible to ensure that they cut the cost of production to a maximum of 30% hence, the investment will not be viable and might not be recommended for investment especially if the investment will be finance by bank loan.

At the calculated revenue, the rate of return was 224.95% but when the revenue was reduced by 10%, rate of return dropped to 192.45%, at 30% drop in revenue, the rate of return dropped to 127.46%. therefore, for the enterprise to remain profitable, the decrease in revenue should not go beyond 30%.

TABLE 9: Sensitivity analysis of Rate of Return on Investment of bio-fortified cassava (Increasing Cost)

VARIABLE RORI	COST	RETURN	RORI	REMARK
Actual cost	87448.32	284159.27	224.95%	Actual estimate
+10% cost	96193.15	284159.27	195.40%	Recommended
+15% cost	100565.56	284159.27	182.56%	Recommended
+20% cost	104937.98	284159.27	170.78%	Recommended
+25% cost	109310.40	284159.27	159.95%	Recommended
+30% cost	113682.81	284159.27	149.95%	Recommended
+35% cost	118055.23	284159.27	140.70%	Not Recommended

Source: Data analysis, 2018

TABLE 10: Sensitivity analysis of Rate of Return on Investment of bio-fortified cassava
(Decreasing revenue)

VARIABLE RORI	COST	RETURN	RORI	REMARK
Actual cost	87448.32	284159.27	224.95%	Actual estimate
-10% revenue	87448.32	255743.34	192.45%	Recommended
-15% revenue	87448.32	241535.37	176.20%	Recommended
-20% revenue	87448.32	227327.41	159.95%	Recommended
-25% revenue	87448.32	213119.45	143.70%	Recommended
-30% revenue	87448.32	198911.48	127.46%	Not Recommended
-35% revenue	87448.32	184703.52	111.21%	Not Recommended

Source: Data analysis, 2018

CONCLUSION AND RECOMMENDATIONS

The study examined the profitability of bio-fortified cassava production and projected the future farm size of the production of bio-fortified cassava in Oyo State, Nigeria. The study concluded that producers of bio-fortified cassava were young and active. The study further found that production of bio-fortified cassava is a profitable enterprise in the study area. The study showed that bio-fortified cassava production can adjust positively to incidentals such as general price inflation and price changes for inputs and outputs that may occur in time. The result of the study concluded that bio-fortified cassava farmers in the study area were small scale farmers but they could increase their production in the future if certain measures were put in place. The bio-fortified cassava farmers have a great potential to boost production through increases in farm sizes of the bio-fortified cassava famers until the year 2026 when equilibrium would be attained at about 2.85ha. in other to adequately achieve these goals, more improved varieties of bio-fortified cassava should be provided, and also, infrastructures should be put in place to help boost farmers moral in their cause of production in the study area.

ACKNOWLEDGEMNT

The authors will like to appreciate the effort of ADPs and Cassava producers' association for the help rendered during the time of data collection.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Alimi, T., O.I. Baruwa, J.O. Bifarin, P.O. Abogan and O.C. Ajewole, (2007). Application of Markov Chain in Forecasting Plantain Farm Size in the Rain Forest Zone of Osun State, Nigeria. *Journal of Agricultural and Rural Development*, 2 (1), 69-83.
- Amina, F.O. Akhigbe-Ahonkai, C. E. (2017). Profitability and Technical Efficiency of Grasscutter production in Nigeria: The case of Ekiti State. *Agricultural Tropica et Subtropica*, 50(1),27–35.
- Anders, H. (2016). Economies of Size and Scale in Agriculture: an interpretive review of empirical measurement. *Review of agricultural economics*, 3(1), 227-238.
- Baruwa, I.O.I Masuku, M.B. and Alimi, I.T. (2011). Economic Analysis of Plantain Production in Derived Savannah Zone of Osun State, Nigeria. *Asian Journal of Agricultural Sciences*, 3(5), 401-407.
- Dachin, Anca, Ursu and Ana (2016). Sensitivity of Gross Margin for Field Crops, in Agrarian Economy and Rural Development - Realities and Perspectives for Romania. The Research Institute for Agricultural Economy and Rural Development (ICEADR). *Bucharest*. 142. Available at: <http://hdl.handle.net/10419/163365>.
- FAO, 2003, Food and Agriculture Organization (FAO) website. www.fao.org.
- Hillocks R., (2002). “Cassava in Africa”. CABI Publishing.
- IITA (2007). A strategy for industrialization of cassava in Africa: Proceedings of a small Group meeting, 14-18 November 2005, Ibadan, Nigeria.
- Iloa, P., Bouis, H. E., Palenberg, M., Moursi, M., Oparinde, A. (2017). Vitamin A Cassava in Nigeria: Crop Development and Delivery. *Afr. J. Food Agric. Nutr. Dev.*, 17(2), 12000–12025.
- NPC (National Population Commission). (2006). Population and Housing Census. Population Local Government Area and Sex. Retrieved from: <http://population.gov.ng/coreactivities/surveys/dataset/2006-phc-priority-tables/>
- Ogunleye Ayodeji Sunday, Bamire Adebayo Simeon, Awolola Olatundun.(2019). Profitability of Investment and Farm Level Efficiency Among Groups of Vitamin A Cassava Farmers in Oyo State Nigeria. *American Journal of Environmental and Resource Economics*, 8(1),14-19. doi: 10.11648/j.eco.20190801.13
- Olatidoye, M. S. Kehinde, A. D. and Alimi, T. (2018). Forecasting the Future Farm Size in Grass Cutter Production in Osun State, Nigeria: A Markov Chain Approach. *Asian Journal of Research in Agriculture and Forestry*. 2(3), 1-10.
- Oparinde, A., Abdoulaye, T., Mignouna, D. B., Bamire, A. S. (2017). Will farmers intend to cultivate Provitamin A genetically modified (GM) cassava in Nigeria? Evidence from a k-means segmentation analysis of beliefs and attitudes. *PLoS ONE* 12(7), e0179427.
- Otekunrin, O.A and Sawicka, B. (2019). “Cassava, a 21st century staple crop: How can Nigeria harness its Enormous Trade potentials?” *Acta Scientific Agriculture* 3(8), 194-202.
- Pannell, D.J. (1997). Sensitivity analysis of normative economic models: Theoretical framework and Practical Strategies. *Journal of Agricultural Economics*. 16, 139-152.
- Phillips T. P., Taylor D. S., Sanni L. & Akoroda, M.O., (2004). A cassava industrial revolution in Nigeria: The Potential for a new industrial crop”. International Institute of Tropical

Agriculture, Ibadan, Nigeria”. International Fund for Agricultural Development, Food and Agriculture Organization of the United Nations, Rome, Italy.

Saltzman, A., Birol, E., Bouis, H. E., Boy, E., De Moura, F. F., Islam, Y. Pfeiffer, W. H. (2014). Biofortification: Progress towards a more nourishing future. Bread and Brain, Education and Poverty. Pontifical Academy of Sciences, Vatican City. Scripta varia 125. Retrieved from: www.pas.va/content/dam/accademia/pdf/sv125/sv125-bouis.pdf